

表 1508 降圧薬使用の有無 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン 濃度	なし	あり	合計	不明	記載なし	総計
5未満	59	180	239		22	261
(%)	(24.7)	(75.3)	(100.0)			
5～	87	237	324		38	362
(%)	(26.9)	(73.1)	(100.0)			
6～	325	471	796	6	140	942
(%)	(40.8)	(59.2)	(100.0)			
7～	1,161	1,927	3,088	22	468	3,578
(%)	(37.6)	(62.4)	(100.0)			
8～	3,686	6,578	10,264	54	1,408	11,726
(%)	(35.9)	(64.1)	(100.0)			
9～	10,303	22,062	32,365	147	4,526	37,038
(%)	(31.8)	(68.2)	(100.0)			
10～	20,514	45,212	65,726	266	9,225	75,217
(%)	(31.2)	(68.8)	(100.0)			
11～	17,028	36,799	53,827	243	7,409	61,479
(%)	(31.6)	(68.4)	(100.0)			
12～	6,596	13,043	19,639	97	2,634	22,370
(%)	(33.6)	(66.4)	(100.0)			
13～	1,837	2,992	4,829	23	685	5,537
(%)	(38.0)	(62.0)	(100.0)			
14～	745	970	1,715	6	212	1,933
(%)	(43.4)	(56.6)	(100.0)			
合計	62,341	130,471	192,812	864	26,767	220,443
(%)	(32.3)	(67.7)	(100.0)			
記載なし	1,025	2,165	3,190	59	5,279	8,528
(%)	(32.1)	(67.9)	(100.0)			
総計	63,366	132,636	196,002	923	32,046	228,971
(%)	(32.3)	(67.7)	(100.0)			
平均	10.70	10.69	10.69	10.67	10.67	10.69
標準偏差	1.34	1.25	1.28	1.32	1.27	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査