

表 1510 降圧薬使用の有無 HDL-C濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

HDL-C濃度	なし	あり	合計	不明	記載なし	総計
20未満	265	328	593	1	77	671
(%)	(44.7)	(55.3)	(100.0)			
20～	3,747	7,646	11,393	24	1,429	12,846
(%)	(32.9)	(67.1)	(100.0)			
30～	10,448	22,232	32,680	67	4,141	36,888
(%)	(32.0)	(68.0)	(100.0)			
40～	11,665	24,422	36,087	82	4,560	40,729
(%)	(32.3)	(67.7)	(100.0)			
50～	8,083	17,573	25,656	56	3,368	29,080
(%)	(31.5)	(68.5)	(100.0)			
60～	4,519	10,254	14,773	38	1,981	16,792
(%)	(30.6)	(69.4)	(100.0)			
70～	2,273	4,886	7,159	22	1,020	8,201
(%)	(31.8)	(68.2)	(100.0)			
80～	1,041	2,276	3,317	2	515	3,834
(%)	(31.4)	(68.6)	(100.0)			
90～	453	1,026	1,479	6	248	1,733
(%)	(30.6)	(69.4)	(100.0)			
100～	375	819	1,194		264	1,458
(%)	(31.4)	(68.6)	(100.0)			
合計	42,869	91,462	134,331	298	17,603	152,232
(%)	(31.9)	(68.1)	(100.0)			
記載なし	20,497	41,174	61,671	625	14,443	76,739
(%)	(33.2)	(66.8)	(100.0)			
総計	63,366	132,636	196,002	923	32,046	228,971
(%)	(32.3)	(67.7)	(100.0)			
平均	47.86	48.23	48.11	48.37	49.33	48.25
標準偏差	16.44	16.34	16.37	15.43	17.75	16.54

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査