

表 1514 降圧薬使用の有無 透析前脈拍(拍/分)別(施設血液透析患者 透析回数週3回)

透析前脈拍	なし	あり	合計	不明	記載なし	総計
40未満	53	82	135	1	19	155
(%)	(39.3)	(60.7)	(100.0)			
40～	481	1,247	1,728	4	197	1,929
(%)	(27.8)	(72.2)	(100.0)			
50～	3,955	10,086	14,041	73	1,659	15,773
(%)	(28.2)	(71.8)	(100.0)			
60～	13,168	30,159	43,327	173	5,241	48,741
(%)	(30.4)	(69.6)	(100.0)			
70～	16,084	35,452	51,536	240	6,408	58,184
(%)	(31.2)	(68.8)	(100.0)			
80～	10,884	21,394	32,278	159	4,286	36,723
(%)	(33.7)	(66.3)	(100.0)			
90～	4,971	9,136	14,107	53	1,758	15,918
(%)	(35.2)	(64.8)	(100.0)			
100～	1,727	2,609	4,336	14	554	4,904
(%)	(39.8)	(60.2)	(100.0)			
110～	440	586	1,026	5	126	1,157
(%)	(42.9)	(57.1)	(100.0)			
120～	201	253	454	1	43	498
(%)	(44.3)	(55.7)	(100.0)			
合計	51,964	111,004	162,968	723	20,291	183,982
(%)	(31.9)	(68.1)	(100.0)			
記載なし	11,402	21,632	33,034	200	11,755	44,989
(%)	(34.5)	(65.5)	(100.0)			
総計	63,366	132,636	196,002	923	32,046	228,971
(%)	(32.3)	(67.7)	(100.0)			
平均	75.46	73.99	74.46	74.41	74.81	74.50
標準偏差	13.20	12.61	12.82	12.38	12.63	12.80

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査