

表 1518 降圧薬使用の有無 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	5,160	7,920	13,080	33	1,963	15,076
(%)	(39.4)	(60.6)	(100.0)			
慢性腎盂腎炎	155	211	366	3	53	422
(%)	(42.3)	(57.7)	(100.0)			
急速進行性糸球体腎炎	86	149	235		49	284
(%)	(36.6)	(63.4)	(100.0)			
妊娠腎/妊娠中毒症	127	173	300	2	57	359
(%)	(42.3)	(57.7)	(100.0)			
その他の分類不能の腎炎	66	94	160	1	34	195
(%)	(41.3)	(58.8)	(100.0)			
多発性嚢胞腎	478	769	1,247	2	215	1,464
(%)	(38.3)	(61.7)	(100.0)			
腎硬化症	778	1,722	2,500	4	382	2,886
(%)	(31.1)	(68.9)	(100.0)			
悪性高血圧	65	202	267	1	44	312
(%)	(24.3)	(75.7)	(100.0)			
糖尿病性腎症	2,962	8,365	11,327	23	1,823	13,173
(%)	(26.1)	(73.9)	(100.0)			
SLE腎炎	133	191	324	1	38	363
(%)	(41.0)	(59.0)	(100.0)			
アミロイド腎	18	33	51		8	59
(%)	(35.3)	(64.7)	(100.0)			
痛風腎	49	88	137	2	15	154
(%)	(35.8)	(64.2)	(100.0)			
先天性代謝異常による腎不全	14	23	37		7	44
(%)	(37.8)	(62.2)	(100.0)			
腎・尿路結核	15	4	19		3	22
(%)	(78.9)	(21.1)	(100.0)			
腎・尿路結石	19	42	61		8	69
(%)	(31.1)	(68.9)	(100.0)			
腎・尿路腫瘍	28	48	76		9	85
(%)	(36.8)	(63.2)	(100.0)			
閉塞性尿路障害	30	32	62		12	74
(%)	(48.4)	(51.6)	(100.0)			
骨髄腫	12	6	18		1	19
(%)	(66.7)	(33.3)	(100.0)			
腎形成不全	47	40	87	2	12	101
(%)	(54.0)	(46.0)	(100.0)			
不明	1,065	1,849	2,914	18	548	3,480
(%)	(36.5)	(63.5)	(100.0)			
再導入	147	223	370	2	44	416
(%)	(39.7)	(60.3)	(100.0)			
その他	294	435	729	2	158	889
(%)	(40.3)	(59.7)	(100.0)			
合計	11,748	22,619	34,367	96	5,483	39,946
(%)	(34.2)	(65.8)	(100.0)			
記載なし						
(%)						
総計	11,748	22,619	34,367	96	5,483	39,946
(%)	(34.2)	(65.8)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査