

表 1523 降圧薬使用の有無 透析時間(時間/回)別 (血液透析濾過患者 透析回数週3回)

透析時間	なし	あり	合計	不明	記載なし	総計
3.0未満	35	50	85		7	92
(%)	(41.2)	(58.8)	(100.0)			
3.0～	655	1,267	1,922	2	358	2,282
(%)	(34.1)	(65.9)	(100.0)			
3.5～	712	1,442	2,154	5	320	2,479
(%)	(33.1)	(66.9)	(100.0)			
4.0～	7,582	15,104	22,686	79	3,669	26,434
(%)	(33.4)	(66.6)	(100.0)			
4.5～	1,209	2,185	3,394	7	447	3,848
(%)	(35.6)	(64.4)	(100.0)			
5.0～	1,311	2,259	3,570	2	460	4,032
(%)	(36.7)	(63.3)	(100.0)			
5.5～	85	119	204		14	218
(%)	(41.7)	(58.3)	(100.0)			
6.0～	117	145	262		22	284
(%)	(44.7)	(55.3)	(100.0)			
6.5～	7	9	16		1	17
(%)	(43.8)	(56.3)	(100.0)			
7.0～	10	23	33		3	36
(%)	(30.3)	(69.7)	(100.0)			
合計	11,723	22,603	34,326	95	5,301	39,722
(%)	(34.2)	(65.8)	(100.0)			
記載なし	25	16	41	1	182	224
(%)	(61.0)	(39.0)	(100.0)			
総計	11,748	22,619	34,367	96	5,483	39,946
(%)	(34.2)	(65.8)	(100.0)			
平均	4.12	4.10	4.10	4.02	4.07	4.10
標準偏差	0.53	0.50	0.51	0.27	0.47	0.50

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査