

表 1525 降圧薬使用の有無 身長(cm)別（血液透析濾過患者 透析回数週3回）

身長	なし	あり	合計	不明	記載なし	総計
130未満	15	20	35		3	38
(%)	(42.9)	(57.1)	(100.0)			
130～	27	32	59	1	10	70
(%)	(45.8)	(54.2)	(100.0)			
135～	128	134	262	1	26	289
(%)	(48.9)	(51.1)	(100.0)			
140～	400	637	1,037	6	120	1,163
(%)	(38.6)	(61.4)	(100.0)			
145～	995	1,626	2,621	11	363	2,995
(%)	(38.0)	(62.0)	(100.0)			
150～	1,733	2,928	4,661	9	542	5,212
(%)	(37.2)	(62.8)	(100.0)			
155～	1,973	3,442	5,415	9	655	6,079
(%)	(36.4)	(63.6)	(100.0)			
160～	2,169	4,248	6,417	17	743	7,177
(%)	(33.8)	(66.2)	(100.0)			
165～	1,889	4,107	5,996	16	710	6,722
(%)	(31.5)	(68.5)	(100.0)			
170～	1,337	2,930	4,267	11	507	4,785
(%)	(31.3)	(68.7)	(100.0)			
175～	488	1,093	1,581	5	200	1,786
(%)	(30.9)	(69.1)	(100.0)			
180～	109	336	445		48	493
(%)	(24.5)	(75.5)	(100.0)			
185～	18	52	70		7	77
(%)	(25.7)	(74.3)	(100.0)			
190～	2	6	8		1	9
(%)	(25.0)	(75.0)	(100.0)			
合計	11,283	21,591	32,874	86	3,935	36,895
(%)	(34.3)	(65.7)	(100.0)			
記載なし	465	1,028	1,493	10	1,548	3,051
(%)	(31.1)	(68.9)	(100.0)			
総計	11,748	22,619	34,367	96	5,483	39,946
(%)	(34.2)	(65.8)	(100.0)			
平均	159.89	161.20	160.75	159.43	160.64	160.74
標準偏差	9.65	9.45	9.54	10.49	9.49	9.54

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査