

表 1526 降圧薬使用の有無 透析後体重(kg)別（血液透析濾過患者 透析回数週3回）

透析後体重	なし	あり	合計	不明	記載なし	総計
20未満	5	2	7		5	12
(%)	(71.4)	(28.6)	(100.0)			
20～	31	35	66		10	76
(%)	(47.0)	(53.0)	(100.0)			
30～	880	1,450	2,330	8	344	2,682
(%)	(37.8)	(62.2)	(100.0)			
40～	3,263	5,499	8,762	15	1,228	10,005
(%)	(37.2)	(62.8)	(100.0)			
50～	3,684	7,275	10,959	25	1,506	12,490
(%)	(33.6)	(66.4)	(100.0)			
60～	2,392	4,926	7,318	13	1,034	8,365
(%)	(32.7)	(67.3)	(100.0)			
70～	935	2,071	3,006	3	414	3,423
(%)	(31.1)	(68.9)	(100.0)			
80～	337	791	1,128	6	139	1,273
(%)	(29.9)	(70.1)	(100.0)			
90～	106	288	394	1	60	455
(%)	(26.9)	(73.1)	(100.0)			
100～	34	88	122		20	142
(%)	(27.9)	(72.1)	(100.0)			
110～	15	39	54	1	6	61
(%)	(27.8)	(72.2)	(100.0)			
120～	8	27	35		2	37
(%)	(22.9)	(77.1)	(100.0)			
合計	11,690	22,491	34,181	72	4,768	39,021
(%)	(34.2)	(65.8)	(100.0)			
記載なし	58	128	186	24	715	925
(%)	(31.2)	(68.8)	(100.0)			
総計	11,748	22,619	34,367	96	5,483	39,946
(%)	(34.2)	(65.8)	(100.0)			
平均	55.71	57.17	56.67	56.80	56.33	56.63
標準偏差	12.73	13.16	13.03	15.30	12.98	13.03

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査