

表 1529 降圧薬使用の有無 Body Mass Index別（血液透析濾過患者 透析回数週3回）

Body Mass Index	なし	あり	合計	不明	記載なし	総計
14未満	55	80	135		14	149
(%)	(40.7)	(59.3)	(100.0)			
14～	356	549	905	2	139	1,046
(%)	(39.3)	(60.7)	(100.0)			
16～	1,233	2,317	3,550	8	440	3,998
(%)	(34.7)	(65.3)	(100.0)			
18～	2,390	4,473	6,863	10	741	7,614
(%)	(34.8)	(65.2)	(100.0)			
20～	2,643	4,915	7,558	14	890	8,462
(%)	(35.0)	(65.0)	(100.0)			
22～	1,998	3,854	5,852	10	709	6,571
(%)	(34.1)	(65.9)	(100.0)			
24～	1,180	2,489	3,669	5	414	4,088
(%)	(32.2)	(67.8)	(100.0)			
26～	652	1,310	1,962	4	231	2,197
(%)	(33.2)	(66.8)	(100.0)			
28～	345	716	1,061	1	115	1,177
(%)	(32.5)	(67.5)	(100.0)			
30～	303	625	928	4	85	1,017
(%)	(32.7)	(67.3)	(100.0)			
合計	11,155	21,328	32,483	58	3,778	36,319
(%)	(34.3)	(65.7)	(100.0)			
記載なし	593	1,291	1,884	38	1,705	3,627
(%)	(31.5)	(68.5)	(100.0)			
総計	11,748	22,619	34,367	96	5,483	39,946
(%)	(34.2)	(65.8)	(100.0)			
平均	21.56	21.74	21.68	21.75	21.55	21.67
標準偏差	3.63	3.66	3.65	3.97	3.60	3.65

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査