

表 1530 降圧薬使用の有無 透析前BUN(mg/dL)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

透析前BUN	なし	あり	合計	不明	記載なし	総計
20未満	17	16	33		6	39
(%)	(51.5)	(48.5)	(100.0)			
20～	111	162	273	1	35	309
(%)	(40.7)	(59.3)	(100.0)			
30～	510	955	1,465	2	186	1,653
(%)	(34.8)	(65.2)	(100.0)			
40～	1,429	2,752	4,181	10	594	4,785
(%)	(34.2)	(65.8)	(100.0)			
50～	2,532	4,751	7,283	14	1,021	8,318
(%)	(34.8)	(65.2)	(100.0)			
60～	2,698	5,015	7,713	22	1,069	8,804
(%)	(35.0)	(65.0)	(100.0)			
70～	1,825	3,458	5,283	11	792	6,086
(%)	(34.5)	(65.5)	(100.0)			
80～	871	1,611	2,482	1	348	2,831
(%)	(35.1)	(64.9)	(100.0)			
90～	353	565	918	1	132	1,051
(%)	(38.5)	(61.5)	(100.0)			
100～	99	148	247	1	31	279
(%)	(40.1)	(59.9)	(100.0)			
110～	15	48	63		11	74
(%)	(23.8)	(76.2)	(100.0)			
120～	5	14	19		1	20
(%)	(26.3)	(73.7)	(100.0)			
合計	10,465	19,495	29,960	63	4,226	34,249
(%)	(34.9)	(65.1)	(100.0)			
記載なし	39	100	139	24	559	722
(%)	(28.1)	(71.9)	(100.0)			
総計	10,504	19,595	30,099	87	4,785	34,971
(%)	(34.9)	(65.1)	(100.0)			
平均	62.28	62.17	62.21	60.56	62.41	62.23
標準偏差	15.22	15.04	15.11	13.56	14.95	15.08

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査