

表 1537 降圧薬使用の有無 透析前CRP濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

透析前CRP濃度	なし	あり	合計	不明	記載なし	総計
0.2未満	5,725	12,661	18,386	35	2,283	20,704
(%)	(31.1)	(68.9)	(100.0)			
0.2～	2,080	3,636	5,716	15	746	6,477
(%)	(36.4)	(63.6)	(100.0)			
0.5～	1,030	1,742	2,772	5	396	3,173
(%)	(37.2)	(62.8)	(100.0)			
1.0～	669	1,061	1,730	6	267	2,003
(%)	(38.7)	(61.3)	(100.0)			
2.0～	412	580	992	4	138	1,134
(%)	(41.5)	(58.5)	(100.0)			
4.0～	153	205	358	1	48	407
(%)	(42.7)	(57.3)	(100.0)			
6.0～	57	91	148	2	21	171
(%)	(38.5)	(61.5)	(100.0)			
8.0～	37	38	75		19	94
(%)	(49.3)	(50.7)	(100.0)			
10.0～	39	56	95		13	108
(%)	(41.1)	(58.9)	(100.0)			
15.0～	27	47	74	2	8	84
(%)	(36.5)	(63.5)	(100.0)			
合計	10,229	20,117	30,346	70	3,939	34,355
(%)	(33.7)	(66.3)	(100.0)			
記載なし	1,519	2,502	4,021	26	1,544	5,591
(%)	(37.8)	(62.2)	(100.0)			
総計	11,748	22,619	34,367	96	5,483	39,946
(%)	(34.2)	(65.8)	(100.0)			
平均	0.63	0.51	0.55	1.22	0.58	0.55
標準偏差	1.73	1.79	1.77	3.53	1.63	1.76

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査