

表 1539 降圧薬使用の有無 透析前リン濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

透析前リン濃度	なし	あり	合計	不明	記載なし	総計
3未満	442	569	1,011	3	119	1,133
(%)	(43.7)	(56.3)	(100.0)			
3.0～	512	861	1,373	3	198	1,574
(%)	(37.3)	(62.7)	(100.0)			
3.5～	866	1,640	2,506	4	339	2,849
(%)	(34.6)	(65.4)	(100.0)			
4.0～	1,376	2,598	3,974	12	579	4,565
(%)	(34.6)	(65.4)	(100.0)			
4.5～	1,696	3,344	5,040	11	661	5,712
(%)	(33.7)	(66.3)	(100.0)			
5.0～	1,821	3,570	5,391	11	732	6,134
(%)	(33.8)	(66.2)	(100.0)			
5.5～	1,582	3,057	4,639	7	653	5,299
(%)	(34.1)	(65.9)	(100.0)			
6.0～	330	519	849	1	100	950
(%)	(38.9)	(61.1)	(100.0)			
6.1～	897	1,911	2,808	4	432	3,244
(%)	(31.9)	(68.1)	(100.0)			
6.5～	819	1,646	2,465	5	336	2,806
(%)	(33.2)	(66.8)	(100.0)			
7.0～	519	1,047	1,566	3	206	1,775
(%)	(33.1)	(66.9)	(100.0)			
7.5～	327	655	982	1	121	1,104
(%)	(33.3)	(66.7)	(100.0)			
8.0～	191	394	585	1	65	651
(%)	(32.6)	(67.4)	(100.0)			
8.5～	116	258	374	2	47	423
(%)	(31.0)	(69.0)	(100.0)			
9.0～	183	410	593	2	87	682
(%)	(30.9)	(69.1)	(100.0)			
合計	11,677	22,479	34,156	70	4,675	38,901
(%)	(34.2)	(65.8)	(100.0)			
記載なし	71	140	211	26	808	1,045
(%)	(33.6)	(66.4)	(100.0)			
総計	11,748	22,619	34,367	96	5,483	39,946
(%)	(34.2)	(65.8)	(100.0)			
平均	5.30	5.39	5.36	5.31	5.35	5.36
標準偏差	1.43	1.42	1.43	1.51	1.42	1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査