

表 1540 降圧薬使用の有無 PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:intact-PTH

PTH値	なし	あり	合計	不明	記載なし	総計
20未満	584	900	1,484	4	195	1,683
(%)	(39.4)	(60.6)	(100.0)			
20～	672	1,251	1,923	3	269	2,195
(%)	(34.9)	(65.1)	(100.0)			
40～	709	1,435	2,144	1	292	2,437
(%)	(33.1)	(66.9)	(100.0)			
60～	787	1,504	2,291	8	293	2,592
(%)	(34.4)	(65.6)	(100.0)			
80～	856	1,656	2,512	5	329	2,846
(%)	(34.1)	(65.9)	(100.0)			
100～	833	1,604	2,437	4	354	2,795
(%)	(34.2)	(65.8)	(100.0)			
120～	752	1,527	2,279	4	298	2,581
(%)	(33.0)	(67.0)	(100.0)			
140～	625	1,419	2,044	8	271	2,323
(%)	(30.6)	(69.4)	(100.0)			
160～	605	1,271	1,876	4	270	2,150
(%)	(32.2)	(67.8)	(100.0)			
180～	1,456	2,914	4,370	5	630	5,005
(%)	(33.3)	(66.7)	(100.0)			
240～	23	46	69		8	77
(%)	(33.3)	(66.7)	(100.0)			
241～	873	1,737	2,610	2	343	2,955
(%)	(33.4)	(66.6)	(100.0)			
300～	506	998	1,504	4	206	1,714
(%)	(33.6)	(66.4)	(100.0)			
360～	310	596	906	1	135	1,042
(%)	(34.2)	(65.8)	(100.0)			
420～	199	302	501	2	88	591
(%)	(39.7)	(60.3)	(100.0)			
480～	47	85	132		14	146
(%)	(35.6)	(64.4)	(100.0)			
500～	77	121	198		39	237
(%)	(38.9)	(61.1)	(100.0)			
540～	81	131	212	1	39	252
(%)	(38.2)	(61.8)	(100.0)			
600～	145	223	368	1	54	423
(%)	(39.4)	(60.6)	(100.0)			
800～	124	185	309	2	39	350
(%)	(40.1)	(59.9)	(100.0)			
合計	10,264	19,905	30,169	59	4,166	34,394
(%)	(34.0)	(66.0)	(100.0)			
記載なし	202	316	518		216	734
(%)	(39.0)	(61.0)	(100.0)			
総計	10,466	20,221	30,687	59	4,382	35,128
(%)	(34.1)	(65.9)	(100.0)			
平均	180.99	174.84	176.93	189.08	179.44	177.26
標準偏差	189.05	160.45	170.74	193.37	159.46	169.45

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査