

表 1541 降圧薬使用の有無 PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	なし	あり	合計	不明	記載なし	総計
20未満	86	149	235		14	249
(%)	(36.6)	(63.4)	(100.0)			
20～	97	181	278	1	22	301
(%)	(34.9)	(65.1)	(100.0)			
35～	80	174	254	1	19	274
(%)	(31.5)	(68.5)	(100.0)			
50～	125	284	409		35	444
(%)	(30.6)	(69.4)	(100.0)			
75～	108	261	369	2	20	391
(%)	(29.3)	(70.7)	(100.0)			
100～	89	225	314		17	331
(%)	(28.3)	(71.7)	(100.0)			
125～	65	151	216	1	17	234
(%)	(30.1)	(69.9)	(100.0)			
150～		4	4			4
(%)		(100.0)	(100.0)			
151～	66	137	203		12	215
(%)	(32.5)	(67.5)	(100.0)			
180～	46	81	127		10	137
(%)	(36.2)	(63.8)	(100.0)			
210～	34	57	91		7	98
(%)	(37.4)	(62.6)	(100.0)			
240～	32	48	80		8	88
(%)	(40.0)	(60.0)	(100.0)			
300～	10	27	37		5	42
(%)	(27.0)	(73.0)	(100.0)			
350～	12	17	29			29
(%)	(41.4)	(58.6)	(100.0)			
400～	4	17	21		3	24
(%)	(19.0)	(81.0)	(100.0)			
500～	17	26	43		2	45
(%)	(39.5)	(60.5)	(100.0)			
合計	871	1,839	2,710	5	191	2,906
(%)	(32.1)	(67.9)	(100.0)			
記載なし	3	5	8		23	31
(%)	(37.5)	(62.5)	(100.0)			
総計	874	1,844	2,718	5	214	2,937
(%)	(32.2)	(67.8)	(100.0)			
平均	114.09	110.75	111.82	71.20	112.37	111.79
標準偏差	116.46	104.69	108.60	43.85	102.37	108.12

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査