

表 1542 降圧薬使用の有無 透析前ヘモグロビン濃度(g/dL)別（血液透析濾過患者 透析回数週3回）

透析前ヘモグロビン 濃度	なし	あり	合計	不明	記載なし	総計
5未満	9	15	24		2	26
(%)	(37.5)	(62.5)	(100.0)			
5～	7	23	30		1	31
(%)	(23.3)	(76.7)	(100.0)			
6～	52	55	107		7	114
(%)	(48.6)	(51.4)	(100.0)			
7～	174	232	406	1	38	445
(%)	(42.9)	(57.1)	(100.0)			
8～	526	910	1,436	5	196	1,637
(%)	(36.6)	(63.4)	(100.0)			
9～	1,700	3,427	5,127	11	712	5,850
(%)	(33.2)	(66.8)	(100.0)			
10～	3,595	7,728	11,323	26	1,598	12,947
(%)	(31.7)	(68.3)	(100.0)			
11～	3,464	6,682	10,146	17	1,414	11,577
(%)	(34.1)	(65.9)	(100.0)			
12～	1,416	2,442	3,858	9	515	4,382
(%)	(36.7)	(63.3)	(100.0)			
13～	449	600	1,049	1	135	1,185
(%)	(42.8)	(57.2)	(100.0)			
14～	219	250	469	1	53	523
(%)	(46.7)	(53.3)	(100.0)			
合計	11,611	22,364	33,975	71	4,671	38,717
(%)	(34.2)	(65.8)	(100.0)			
記載なし	137	255	392	25	812	1,229
(%)	(34.9)	(65.1)	(100.0)			
総計	11,748	22,619	34,367	96	5,483	39,946
(%)	(34.2)	(65.8)	(100.0)			
平均	10.88	10.82	10.84	10.74	10.84	10.84
標準偏差	1.36	1.23	1.27	1.26	1.19	1.26

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査