

表 1544 降圧薬使用の有無 HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

HDL-C濃度	なし	あり	合計	不明	記載なし	総計
20未満	25	44	69		12	81
(%)	(36.2)	(63.8)	(100.0)			
20～	518	1,130	1,648	1	225	1,874
(%)	(31.4)	(68.6)	(100.0)			
30～	1,721	3,666	5,387	9	735	6,131
(%)	(31.9)	(68.1)	(100.0)			
40～	2,181	4,337	6,518	8	834	7,360
(%)	(33.5)	(66.5)	(100.0)			
50～	1,833	3,417	5,250	6	715	5,971
(%)	(34.9)	(65.1)	(100.0)			
60～	1,181	2,231	3,412	6	439	3,857
(%)	(34.6)	(65.4)	(100.0)			
70～	608	1,142	1,750	3	270	2,023
(%)	(34.7)	(65.3)	(100.0)			
80～	319	567	886	1	126	1,013
(%)	(36.0)	(64.0)	(100.0)			
90～	130	248	378	1	52	431
(%)	(34.4)	(65.6)	(100.0)			
100～	114	213	327		56	383
(%)	(34.9)	(65.1)	(100.0)			
合計	8,630	16,995	25,625	35	3,464	29,124
(%)	(33.7)	(66.3)	(100.0)			
記載なし	3,118	5,624	8,742	61	2,019	10,822
(%)	(35.7)	(64.3)	(100.0)			
総計	11,748	22,619	34,367	96	5,483	39,946
(%)	(34.2)	(65.8)	(100.0)			
平均	51.28	50.54	50.79	52.26	51.33	50.85
標準偏差	17.02	17.12	17.09	15.91	17.85	17.18

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査