

表 1566 降圧薬使用の有無 PTH値(pg/mL)別（腹膜透析患者）PTH測定法:whole-PTH

PTH値	なし	あり	合計	不明	記載なし	総計
20未満	9	31	40			40
(%)	(22.5)	(77.5)	(100.0)			
20～	10	35	45			45
(%)	(22.2)	(77.8)	(100.0)			
35～	12	42	54		1	55
(%)	(22.2)	(77.8)	(100.0)			
50～	22	54	76		1	77
(%)	(28.9)	(71.1)	(100.0)			
75～	21	48	69			69
(%)	(30.4)	(69.6)	(100.0)			
100～	15	56	71			71
(%)	(21.1)	(78.9)	(100.0)			
125～	6	38	44		1	45
(%)	(13.6)	(86.4)	(100.0)			
150～		1	1			1
(%)		(100.0)	(100.0)			
151～	11	31	42		1	43
(%)	(26.2)	(73.8)	(100.0)			
180～	7	25	32		2	34
(%)	(21.9)	(78.1)	(100.0)			
210～	4	23	27			27
(%)	(14.8)	(85.2)	(100.0)			
240～	5	18	23			23
(%)	(21.7)	(78.3)	(100.0)			
300～	1	18	19			19
(%)	(5.3)	(94.7)	(100.0)			
350～	1	7	8			8
(%)	(12.5)	(87.5)	(100.0)			
400～	4	7	11			11
(%)	(36.4)	(63.6)	(100.0)			
500～	2	9	11			11
(%)	(18.2)	(81.8)	(100.0)			
合計	130	443	573		6	579
(%)	(22.7)	(77.3)	(100.0)			
記載なし	52	60	112	1	27	140
(%)	(46.4)	(53.6)	(100.0)			
総計	182	503	685	1	33	719
(%)	(26.6)	(73.4)	(100.0)			
平均	122.34	135.52	132.53		134.83	132.56
標準偏差	128.72	130.27	129.93		65.46	129.39

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査