

表 1585 喫煙の有無 身長(cm)別 (透析患者全体)

身長	現在吸っていない	現在吸っている	合計	不明	記載なし	総計
130未満	285	13	298	20	36	354
(%)	(95.6)	(4.4)	(100.0)			
130～	553	7	560	38	98	696
(%)	(98.8)	(1.3)	(100.0)			
135～	2,274	58	2,332	118	387	2,837
(%)	(97.5)	(2.5)	(100.0)			
140～	8,097	216	8,313	447	1,537	10,297
(%)	(97.4)	(2.6)	(100.0)			
145～	18,134	732	18,866	1,096	3,711	23,673
(%)	(96.1)	(3.9)	(100.0)			
150～	28,392	1,897	30,289	1,844	6,102	38,235
(%)	(93.7)	(6.3)	(100.0)			
155～	31,839	3,309	35,148	2,334	6,990	44,472
(%)	(90.6)	(9.4)	(100.0)			
160～	36,475	5,723	42,198	2,896	8,556	53,650
(%)	(86.4)	(13.6)	(100.0)			
165～	31,170	6,490	37,660	2,736	7,592	47,988
(%)	(82.8)	(17.2)	(100.0)			
170～	19,260	4,886	24,146	1,870	5,058	31,074
(%)	(79.8)	(20.2)	(100.0)			
175～	6,871	1,989	8,860	642	1,817	11,319
(%)	(77.6)	(22.4)	(100.0)			
180～	1,698	630	2,328	201	450	2,979
(%)	(72.9)	(27.1)	(100.0)			
185～	267	105	372	26	67	465
(%)	(71.8)	(28.2)	(100.0)			
190～	42	15	57	6	14	77
(%)	(73.7)	(26.3)	(100.0)			
合計	185,357	26,070	211,427	14,274	42,415	268,116
(%)	(87.7)	(12.3)	(100.0)			
記載なし	9,423	806	10,229	2,555	29,208	41,992
(%)	(92.1)	(7.9)	(100.0)			
総計	194,780	26,876	221,656	16,829	71,623	310,108
(%)	(87.9)	(12.1)	(100.0)			
平均	159.30	164.70	159.96	160.90	160.17	160.05
標準偏差	9.68	8.29	9.68	9.47	9.55	9.65

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査