

表 1739 腹膜透析の経験 透析前リン濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

透析前リン濃度	なし	あり(年末時点でカテーテルあり)	あり(年末時点でカテーテルなし)	合計	不明	記載なし	総計
3未満	7,134	32	144	7,310	45	1,606	8,961
(%)	(97.6)	(0.4)	(2.0)	(100.0)			
3.0～	8,668	37	191	8,896	38	2,046	10,980
(%)	(97.4)	(0.4)	(2.1)	(100.0)			
3.5～	15,060	54	386	15,500	64	3,454	19,018
(%)	(97.2)	(0.3)	(2.5)	(100.0)			
4.0～	22,395	101	564	23,060	96	5,036	28,192
(%)	(97.1)	(0.4)	(2.4)	(100.0)			
4.5～	26,428	116	839	27,383	110	5,975	33,468
(%)	(96.5)	(0.4)	(3.1)	(100.0)			
5.0～	26,419	121	833	27,373	108	6,328	33,809
(%)	(96.5)	(0.4)	(3.0)	(100.0)			
5.5～	22,918	124	727	23,769	83	5,160	29,012
(%)	(96.4)	(0.5)	(3.1)	(100.0)			
6.0～	3,844	23	126	3,993	10	925	4,928
(%)	(96.3)	(0.6)	(3.2)	(100.0)			
6.1～	12,982	67	406	13,455	40	3,010	16,505
(%)	(96.5)	(0.5)	(3.0)	(100.0)			
6.5～	11,228	79	372	11,679	33	2,665	14,377
(%)	(96.1)	(0.7)	(3.2)	(100.0)			
7.0～	6,855	40	233	7,128	22	1,710	8,860
(%)	(96.2)	(0.6)	(3.3)	(100.0)			
7.5～	4,148	24	154	4,326	19	1,014	5,359
(%)	(95.9)	(0.6)	(3.6)	(100.0)			
8.0～	2,489	16	80	2,585	11	596	3,192
(%)	(96.3)	(0.6)	(3.1)	(100.0)			
8.5～	1,536	10	70	1,616	5	402	2,023
(%)	(95.0)	(0.6)	(4.3)	(100.0)			
9.0～	2,453	16	89	2,558	8	590	3,156
(%)	(95.9)	(0.6)	(3.5)	(100.0)			
合計	174,557	860	5,214	180,631	692	40,517	221,840
(%)	(96.6)	(0.5)	(2.9)	(100.0)			
記載なし	3,748	27	101	3,876	33	3,222	7,131
(%)	(96.7)	(0.7)	(2.6)	(100.0)			
総計	178,305	887	5,315	184,507	725	43,739	228,971
(%)	(96.6)	(0.5)	(2.9)	(100.0)			
平均	5.19	5.39	5.36	5.19	5.01	5.22	5.20
標準偏差	1.42	1.49	1.41	1.42	1.45	1.44	1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査