

表 1742 腹膜透析の経験 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

| 透析前ヘモグロビン濃度 | なし      | あり(年末時点でカテーテルあり) | あり(年末時点でカテーテルなし) | 合計      | 不明    | 記載なし   | 総計      |
|-------------|---------|------------------|------------------|---------|-------|--------|---------|
| 5未満         | 227     | 1                | 4                | 232     | 11    | 18     | 261     |
| (%)         | (97.8)  | (0.4)            | (1.7)            | (100.0) |       |        |         |
| 5～          | 312     |                  | 5                | 317     | 3     | 42     | 362     |
| (%)         | (98.4)  |                  | (1.6)            | (100.0) |       |        |         |
| 6～          | 768     | 5                | 14               | 787     | 5     | 150    | 942     |
| (%)         | (97.6)  | (0.6)            | (1.8)            | (100.0) |       |        |         |
| 7～          | 2,857   | 22               | 67               | 2,946   | 20    | 612    | 3,578   |
| (%)         | (97.0)  | (0.7)            | (2.3)            | (100.0) |       |        |         |
| 8～          | 9,425   | 52               | 236              | 9,713   | 58    | 1,955  | 11,726  |
| (%)         | (97.0)  | (0.5)            | (2.4)            | (100.0) |       |        |         |
| 9～          | 29,345  | 152              | 779              | 30,276  | 116   | 6,646  | 37,038  |
| (%)         | (96.9)  | (0.5)            | (2.6)            | (100.0) |       |        |         |
| 10～         | 59,422  | 258              | 1,653            | 61,333  | 212   | 13,672 | 75,217  |
| (%)         | (96.9)  | (0.4)            | (2.7)            | (100.0) |       |        |         |
| 11～         | 48,260  | 244              | 1,576            | 50,080  | 171   | 11,228 | 61,479  |
| (%)         | (96.4)  | (0.5)            | (3.1)            | (100.0) |       |        |         |
| 12～         | 17,515  | 87               | 628              | 18,230  | 78    | 4,062  | 22,370  |
| (%)         | (96.1)  | (0.5)            | (3.4)            | (100.0) |       |        |         |
| 13～         | 4,334   | 34               | 173              | 4,541   | 21    | 975    | 5,537   |
| (%)         | (95.4)  | (0.7)            | (3.8)            | (100.0) |       |        |         |
| 14～         | 1,527   | 9                | 63               | 1,599   | 9     | 325    | 1,933   |
| (%)         | (95.5)  | (0.6)            | (3.9)            | (100.0) |       |        |         |
| 合計          | 173,992 | 864              | 5,198            | 180,054 | 704   | 39,685 | 220,443 |
| (%)         | (96.6)  | (0.5)            | (2.9)            | (100.0) |       |        |         |
| 記載なし        | 4,313   | 23               | 117              | 4,453   | 21    | 4,054  | 8,528   |
| (%)         | (96.9)  | (0.5)            | (2.6)            | (100.0) |       |        |         |
| 総計          | 178,305 | 887              | 5,315            | 184,507 | 725   | 43,739 | 228,971 |
| (%)         | (96.6)  | (0.5)            | (2.9)            | (100.0) |       |        |         |
| 平均          | 10.68   | 10.69            | 10.85            | 10.69   | 10.49 | 10.71  | 10.69   |
| 標準偏差        | 1.28    | 1.38             | 1.28             | 1.28    | 1.62  | 1.24   | 1.28    |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査