

表 1771 腹膜透析の経験 透析前リン濃度(mg/dL)別 (血液透析濾過患者 透析回数週3回)

透析前リン濃度	なし	あり(年末時点でカテーテルあり)	あり(年末時点でカテーテルなし)	合計	不明	記載なし	総計
3未満	890	1	35	926	3	204	1,133
(%)	(96.1)	(0.1)	(3.8)	(100.0)			
3.0～	1,246	5	41	1,292	5	277	1,574
(%)	(96.4)	(0.4)	(3.2)	(100.0)			
3.5～	2,246	12	97	2,355	7	487	2,849
(%)	(95.4)	(0.5)	(4.1)	(100.0)			
4.0～	3,563	11	169	3,743	17	805	4,565
(%)	(95.2)	(0.3)	(4.5)	(100.0)			
4.5～	4,505	18	174	4,697	19	996	5,712
(%)	(95.9)	(0.4)	(3.7)	(100.0)			
5.0～	4,794	29	254	5,077	15	1,042	6,134
(%)	(94.4)	(0.6)	(5.0)	(100.0)			
5.5～	4,171	12	178	4,361	16	922	5,299
(%)	(95.6)	(0.3)	(4.1)	(100.0)			
6.0～	739	4	28	771	2	177	950
(%)	(95.8)	(0.5)	(3.6)	(100.0)			
6.1～	2,514	12	116	2,642	12	590	3,244
(%)	(95.2)	(0.5)	(4.4)	(100.0)			
6.5～	2,204	13	107	2,324	16	466	2,806
(%)	(94.8)	(0.6)	(4.6)	(100.0)			
7.0～	1,386	10	66	1,462	8	305	1,775
(%)	(94.8)	(0.7)	(4.5)	(100.0)			
7.5～	871	3	48	922	4	178	1,104
(%)	(94.5)	(0.3)	(5.2)	(100.0)			
8.0～	509		24	533	2	116	651
(%)	(95.5)		(4.5)	(100.0)			
8.5～	341	3	12	356	2	65	423
(%)	(95.8)	(0.8)	(3.4)	(100.0)			
9.0～	512	8	29	549	3	130	682
(%)	(93.3)	(1.5)	(5.3)	(100.0)			
合計	30,491	141	1,378	32,010	131	6,760	38,901
(%)	(95.3)	(0.4)	(4.3)	(100.0)			
記載なし	296	3	13	312	15	718	1,045
(%)	(94.9)	(1.0)	(4.2)	(100.0)			
総計	30,787	144	1,391	32,322	146	7,478	39,946
(%)	(95.3)	(0.4)	(4.3)	(100.0)			
平均	5.35	5.66	5.44	5.36	5.53	5.35	5.36
標準偏差	1.42	1.59	1.45	1.42	1.51	1.44	1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査