

表 1782 腎移植の回数 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
施設血液透析	187,183	1,821	92	5		189,101	743	65,797	255,641
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
Off-line HDF	4,657	135	7			4,799	19	1,497	6,315
(%)	(97.0)	(2.8)	(0.1)			(100.0)			
On-line HDF	26,378	521	29	2		26,930	130	9,030	36,090
(%)	(98.0)	(1.9)	(0.1)	(0.0)		(100.0)			
Push/Pull HDF	329	4				333		204	537
(%)	(98.8)	(1.2)				(100.0)			
ハイオフィルトレーション	280	5				285		56	341
(%)	(98.2)	(1.8)				(100.0)			
血液濾過	30					30		62	92
(%)	(100.0)					(100.0)			
血液吸着透析 (リクセル)	1,206	70	5	1		1,282	3	338	1,623
(%)	(94.1)	(5.5)	(0.4)	(0.1)		(100.0)			
血液吸着透析 (その他)	5					5		2	7
(%)	(100.0)					(100.0)			
在宅血液透析	326	38	4			368	1	152	521
(%)	(88.6)	(10.3)	(1.1)			(100.0)			
腹膜透析 (CAPD)	3,129	27	1	1		3,158	6	1,857	5,021
(%)	(99.1)	(0.9)	(0.0)	(0.0)		(100.0)			
腹膜透析 (APD)	1,875	17				1,892	13	934	2,839
(%)	(99.1)	(0.9)				(100.0)			
腹膜透析 (CCPD)	771	12	1			784	4	293	1,081
(%)	(98.3)	(1.5)	(0.1)			(100.0)			
合計	226,169	2,650	139	9		228,967	919	80,222	310,108
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
記載なし (%)									
総計	226,169	2,650	139	9		228,967	919	80,222	310,108
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査