

表 1787 腎移植の回数 原疾患別（透析患者全体）

原疾患	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
慢性糸球体腎炎	70,361	981	56	2		71,400	275	25,295	96,970
(%)	(98.5)	(1.4)	(0.1)	(0.0)		(100.0)			
慢性腎盂腎炎	2,215	21	2			2,238	13	791	3,042
(%)	(99.0)	(0.9)	(0.1)			(100.0)			
急速進行性糸球体腎炎	1,825	10	1			1,836	4	654	2,494
(%)	(99.4)	(0.5)	(0.1)			(100.0)			
妊娠腎/妊娠中毒症	1,177	22	1			1,200	4	410	1,614
(%)	(98.1)	(1.8)	(0.1)			(100.0)			
その他の分類不能の腎炎	979	17	1			997	3	373	1,373
(%)	(98.2)	(1.7)	(0.1)			(100.0)			
多発性嚢胞腎	8,288	26	1			8,315	32	2,659	11,006
(%)	(99.7)	(0.3)	(0.0)			(100.0)			
腎硬化症	21,365	42	1			21,408	93	6,797	28,298
(%)	(99.8)	(0.2)	(0.0)			(100.0)			
悪性高血圧	1,866	8				1,874	3	696	2,573
(%)	(99.6)	(0.4)				(100.0)			
糖尿病性腎症	86,893	127	9	1		87,030	346	30,705	118,081
(%)	(99.8)	(0.1)	(0.0)	(0.0)		(100.0)			
SLE腎炎	1,686	16				1,702	8	526	2,236
(%)	(99.1)	(0.9)				(100.0)			
アミロイド腎	336					336	1	114	451
(%)	(100.0)					(100.0)			
痛風腎	845	6	1			852	4	257	1,113
(%)	(99.2)	(0.7)	(0.1)			(100.0)			
先天性代謝異常による腎不全	195	12				207		65	272
(%)	(94.2)	(5.8)				(100.0)			
腎・尿路結核	152					152	1	53	206
(%)	(100.0)					(100.0)			
腎・尿路結石	427					427	3	151	581
(%)	(100.0)					(100.0)			
腎・尿路腫瘍	650	4				654	1	231	886
(%)	(99.4)	(0.6)				(100.0)			
閉塞性尿路障害	550	7				557	2	164	723
(%)	(98.7)	(1.3)				(100.0)			
骨髄腫	214					214		62	276
(%)	(100.0)					(100.0)			
腎形成不全	452	39	2			493	1	155	649
(%)	(91.7)	(7.9)	(0.4)			(100.0)			
不明	19,794	149	7			19,950	102	7,492	27,544
(%)	(99.2)	(0.7)	(0.0)			(100.0)			
再導入	605	1,082	51	6		1,744	1	418	2,163
(%)	(34.7)	(62.0)	(2.9)	(0.3)		(100.0)			
その他	5,294	81	6			5,381	22	2,154	7,557
(%)	(98.4)	(1.5)	(0.1)			(100.0)			
合計	226,169	2,650	139	9		228,967	919	80,222	310,108
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
記載なし									
(%)									
総計	226,169	2,650	139	9		228,967	919	80,222	310,108
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査