

表 1794 腎移植の回数 透析前CRP濃度(mg/dL)別 (透析患者全体)

透析前CRP濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
0.2未満	108,284	1,331	71	3		109,689	372	25,248	135,309
(%)	(98.7)	(1.2)	(0.1)	(0.0)		(100.0)			
0.2～	36,149	420	17	2		36,588	127	8,092	44,807
(%)	(98.8)	(1.1)	(0.0)	(0.0)		(100.0)			
0.5～	18,125	184	11			18,320	72	4,365	22,757
(%)	(98.9)	(1.0)	(0.1)			(100.0)			
1.0～	11,863	135	8			12,006	40	2,702	14,748
(%)	(98.8)	(1.1)	(0.1)			(100.0)			
2.0～	7,180	78	9	1		7,268	32	1,705	9,005
(%)	(98.8)	(1.1)	(0.1)	(0.0)		(100.0)			
4.0～	2,557	31	1			2,589	16	639	3,244
(%)	(98.8)	(1.2)	(0.0)			(100.0)			
6.0～	1,257	6				1,263	8	305	1,576
(%)	(99.5)	(0.5)				(100.0)			
8.0～	684	3		1		688	4	178	870
(%)	(99.4)	(0.4)		(0.1)		(100.0)			
10.0～	825	13	1			839	6	189	1,034
(%)	(98.3)	(1.5)	(0.1)			(100.0)			
15.0～	523	4				527	3	163	693
(%)	(99.2)	(0.8)				(100.0)			
合計	187,447	2,205	118	7		189,777	680	43,586	234,043
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
記載なし	38,722	445	21	2		39,190	239	36,636	76,065
(%)	(98.8)	(1.1)	(0.1)	(0.0)		(100.0)			
総計	226,169	2,650	139	9		228,967	919	80,222	310,108
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
平均	0.63	0.54	0.59	1.75		0.62	0.84	0.66	0.63
標準偏差	1.87	1.47	1.35	3.07		1.86	2.24	2.14	1.92

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査