

表 1795 腎移植の回数 透析前補正カルシウム濃度(mg/dL)別 (透析患者全体)

透析前補正 カルシウム濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
7.0未満	911	24	1			936	1	205	1,142
(%)	(97.3)	(2.6)	(0.1)			(100.0)			
7.0～	1,315	33	2			1,350	1	298	1,649
(%)	(97.4)	(2.4)	(0.1)			(100.0)			
7.5～	4,928	78	5			5,011	11	1,377	6,399
(%)	(98.3)	(1.6)	(0.1)			(100.0)			
8.0～	15,967	195	11	1		16,174	36	4,229	20,439
(%)	(98.7)	(1.2)	(0.1)	(0.0)		(100.0)			
8.4～	4,670	41	2	1		4,714	9	1,149	5,872
(%)	(99.1)	(0.9)	(0.0)	(0.0)		(100.0)			
8.5～	53,058	530	28	2		53,618	197	13,504	67,319
(%)	(99.0)	(1.0)	(0.1)	(0.0)		(100.0)			
9.0～	64,957	749	37	2		65,745	254	16,005	82,004
(%)	(98.8)	(1.1)	(0.1)	(0.0)		(100.0)			
9.5～	42,734	540	25	1		43,300	125	10,544	53,969
(%)	(98.7)	(1.2)	(0.1)	(0.0)		(100.0)			
10.0～	5,268	61	4			5,333	15	1,250	6,598
(%)	(98.8)	(1.1)	(0.1)			(100.0)			
10.1～	13,473	178	12	1		13,664	44	3,269	16,977
(%)	(98.6)	(1.3)	(0.1)	(0.0)		(100.0)			
10.5～	6,501	99	4			6,604	21	1,645	8,270
(%)	(98.4)	(1.5)	(0.1)			(100.0)			
11.0～	2,208	27				2,235	8	538	2,781
(%)	(98.8)	(1.2)				(100.0)			
11.5～	789	10	2			801	7	165	973
(%)	(98.5)	(1.2)	(0.2)			(100.0)			
12.0～	550	5	1			556	4	121	681
(%)	(98.9)	(0.9)	(0.2)			(100.0)			
合計	217,329	2,570	134	8		220,041	733	54,299	275,073
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
記載なし	8,840	80	5	1		8,926	186	25,923	35,035
(%)	(99.0)	(0.9)	(0.1)	(0.0)		(100.0)			
総計	226,169	2,650	139	9		228,967	919	80,222	310,108
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
平均	9.19	9.19	9.20	9.11		9.19	9.25	9.18	9.19
標準偏差	0.74	0.81	0.85	0.66		0.75	0.71	0.74	0.74

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査