

表 1796 腎移植の回数 透析前リン濃度(mg/dL)別 (透析患者全体)

透析前リン濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
3未満	8,659	62	5			8,726	53	2,107	10,886
(%)	(99.2)	(0.7)	(0.1)			(100.0)			
3.0～	10,768	84	4	1		10,857	45	2,680	13,582
(%)	(99.2)	(0.8)	(0.0)	(0.0)		(100.0)			
3.5～	18,824	149	8			18,981	83	4,618	23,682
(%)	(99.2)	(0.8)	(0.0)			(100.0)			
4.0～	28,054	260	11	1		28,326	105	6,851	35,282
(%)	(99.0)	(0.9)	(0.0)	(0.0)		(100.0)			
4.5～	33,329	364	23	2		33,718	123	8,218	42,059
(%)	(98.8)	(1.1)	(0.1)	(0.0)		(100.0)			
5.0～	33,596	388	28			34,012	125	8,581	42,718
(%)	(98.8)	(1.1)	(0.1)			(100.0)			
5.5～	28,998	404	15	1		29,418	99	7,188	36,705
(%)	(98.6)	(1.4)	(0.1)	(0.0)		(100.0)			
6.0～	4,874	63		1		4,938	11	1,312	6,261
(%)	(98.7)	(1.3)		(0.0)		(100.0)			
6.1～	16,586	233	9			16,828	61	4,220	21,109
(%)	(98.6)	(1.4)	(0.1)			(100.0)			
6.5～	14,346	215	13	1		14,575	45	3,658	18,278
(%)	(98.4)	(1.5)	(0.1)	(0.0)		(100.0)			
7.0～	8,740	134	7			8,881	29	2,367	11,277
(%)	(98.4)	(1.5)	(0.1)			(100.0)			
7.5～	5,334	83	6	1		5,424	21	1,413	6,858
(%)	(98.3)	(1.5)	(0.1)	(0.0)		(100.0)			
8.0～	3,191	58	3			3,252	11	790	4,053
(%)	(98.1)	(1.8)	(0.1)			(100.0)			
8.5～	2,007	38	1			2,046	2	524	2,572
(%)	(98.1)	(1.9)	(0.0)			(100.0)			
9.0～	3,156	64	2			3,222	6	849	4,077
(%)	(98.0)	(2.0)	(0.1)			(100.0)			
合計	220,462	2,599	135	8		223,204	819	55,376	279,399
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
記載なし	5,707	51	4	1		5,763	100	24,846	30,709
(%)	(99.0)	(0.9)	(0.1)	(0.0)		(100.0)			
総計	226,169	2,650	139	9		228,967	919	80,222	310,108
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
平均	5.20	5.55	5.41	5.32		5.21	5.00	5.24	5.21
標準偏差	1.42	1.48	1.44	1.43		1.42	1.41	1.44	1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査