

表 1799 腎移植の回数 透析前ヘモグロビン濃度(g/dL)別 (透析患者全体)

透析前ヘモグロビン濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
5未満	270	4	1			275		28	303
(%)	(98.2)	(1.5)	(0.4)			(100.0)			
5～	373	2				375	1	53	429
(%)	(99.5)	(0.5)				(100.0)			
6～	937	9				946	9	205	1,160
(%)	(99.0)	(1.0)				(100.0)			
7～	3,565	44	2	1		3,612	14	809	4,435
(%)	(98.7)	(1.2)	(0.1)	(0.0)		(100.0)			
8～	11,701	126	4			11,831	61	2,662	14,554
(%)	(98.9)	(1.1)	(0.0)			(100.0)			
9～	36,514	373	21			36,908	139	8,942	45,989
(%)	(98.9)	(1.0)	(0.1)			(100.0)			
10～	74,167	802	44	2		75,015	273	18,624	93,912
(%)	(98.9)	(1.1)	(0.1)	(0.0)		(100.0)			
11～	61,542	763	43	4		62,352	213	15,608	78,173
(%)	(98.7)	(1.2)	(0.1)	(0.0)		(100.0)			
12～	22,668	343	15	1		23,027	90	5,737	28,854
(%)	(98.4)	(1.5)	(0.1)	(0.0)		(100.0)			
13～	5,783	97	3			5,883	22	1,403	7,308
(%)	(98.3)	(1.6)	(0.1)			(100.0)			
14～	2,135	29	1			2,165	8	500	2,673
(%)	(98.6)	(1.3)	(0.0)			(100.0)			
合計	219,655	2,592	134	8		222,389	830	54,571	277,790
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
記載なし	6,514	58	5	1		6,578	89	25,651	32,318
(%)	(99.0)	(0.9)	(0.1)	(0.0)		(100.0)			
総計	226,169	2,650	139	9		228,967	919	80,222	310,108
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
平均	10.70	10.85	10.79	10.94		10.71	10.63	10.73	10.71
標準偏差	1.29	1.33	1.28	1.56		1.29	1.34	1.25	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査