

表 1800 腎移植の回数 総コレステロール濃度(mg/dL)別 (透析患者全体)

総コレステロール濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
100未満	6,283	60	3			6,346	32	1,452	7,830
(%)	(99.0)	(0.9)	(0.0)			(100.0)			
100～	17,986	174	15			18,175	84	4,463	22,722
(%)	(99.0)	(1.0)	(0.1)			(100.0)			
120～	33,538	332	18	3		33,891	124	8,186	42,201
(%)	(99.0)	(1.0)	(0.1)	(0.0)		(100.0)			
140～	38,766	466	25	2		39,259	136	9,264	48,659
(%)	(98.7)	(1.2)	(0.1)	(0.0)		(100.0)			
160～	32,071	423	18	1		32,513	117	7,613	40,243
(%)	(98.6)	(1.3)	(0.1)	(0.0)		(100.0)			
180～	20,338	295	14			20,647	47	4,831	25,525
(%)	(98.5)	(1.4)	(0.1)			(100.0)			
200～	10,837	174	9	1		11,021	36	2,666	13,723
(%)	(98.3)	(1.6)	(0.1)	(0.0)		(100.0)			
220～	4,807	70				4,877	15	1,179	6,071
(%)	(98.6)	(1.4)				(100.0)			
240～	1,925	26	3			1,954	5	441	2,400
(%)	(98.5)	(1.3)	(0.2)			(100.0)			
260～	1,221	19	2			1,242	4	340	1,586
(%)	(98.3)	(1.5)	(0.2)			(100.0)			
合計	167,772	2,039	107	7		169,925	600	40,435	210,960
(%)	(98.7)	(1.2)	(0.1)	(0.0)		(100.0)			
記載なし	58,397	611	32	2		59,042	319	39,787	99,148
(%)	(98.9)	(1.0)	(0.1)	(0.0)		(100.0)			
総計	226,169	2,650	139	9		228,967	919	80,222	310,108
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
平均	155.80	161.38	159.20	153.43		155.87	150.89	155.89	155.86
標準偏差	35.68	35.48	43.55	27.71		35.69	35.82	35.94	35.74

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査