

表 1802 腎移植の回数 Non-HDL-C濃度(mg/dL)別（透析患者全体）

Non-HDL-C濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
50未満	2,770	35	1			2,806	7	597	3,410
(%)	(98.7)	(1.2)	(0.0)			(100.0)			
50～	12,656	143	13			12,812	37	3,252	16,101
(%)	(98.8)	(1.1)	(0.1)			(100.0)			
70～	28,059	327	21	2		28,409	85	7,075	35,569
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
90～	33,698	443	18	1		34,160	98	8,120	42,378
(%)	(98.6)	(1.3)	(0.1)	(0.0)		(100.0)			
110～	27,401	339	16	2		27,758	74	6,371	34,203
(%)	(98.7)	(1.2)	(0.1)	(0.0)		(100.0)			
130～	16,760	235	7			17,002	36	3,925	20,963
(%)	(98.6)	(1.4)	(0.0)			(100.0)			
150～	8,451	117	5	1		8,574	20	1,996	10,590
(%)	(98.6)	(1.4)	(0.1)	(0.0)		(100.0)			
170～	3,790	41	5			3,836	12	874	4,722
(%)	(98.8)	(1.1)	(0.1)			(100.0)			
190～	1,452	13				1,465	3	341	1,809
(%)	(99.1)	(0.9)				(100.0)			
210～	857	12	2			871	1	252	1,124
(%)	(98.4)	(1.4)	(0.2)			(100.0)			
合計	135,894	1,705	88	6		137,693	373	32,803	170,869
(%)	(98.7)	(1.2)	(0.1)	(0.0)		(100.0)			
記載なし	90,275	945	51	3		91,274	546	47,419	139,239
(%)	(98.9)	(1.0)	(0.1)	(0.0)		(100.0)			
総計	226,169	2,650	139	9		228,967	919	80,222	310,108
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
平均	107.18	108.52	107.50	106.17		107.19	104.18	106.65	107.08
標準偏差	33.57	33.26	43.28	29.40		33.57	31.53	33.77	33.61

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査