

表 1815 腎移植の回数 体重減少量(kg)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

体重減少量	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
0.0未満	599	3				602	2	151	755
(%)	(99.5)	(0.5)				(100.0)			
0.0～	6,786	46	3			6,835	40	1,575	8,450
(%)	(99.3)	(0.7)	(0.0)			(100.0)			
1.0～	32,714	308	18	1		33,041	143	7,841	41,025
(%)	(99.0)	(0.9)	(0.1)	(0.0)		(100.0)			
2.0～	58,175	629	31	2		58,837	202	14,122	73,161
(%)	(98.9)	(1.1)	(0.1)	(0.0)		(100.0)			
3.0～	30,177	395	19			30,591	92	7,427	38,110
(%)	(98.6)	(1.3)	(0.1)			(100.0)			
4.0～	7,409	82	2			7,493	17	1,935	9,445
(%)	(98.9)	(1.1)	(0.0)			(100.0)			
5.0～	1,532	21				1,553	2	403	1,958
(%)	(98.6)	(1.4)				(100.0)			
6.0～	309	3				312	1	82	395
(%)	(99.0)	(1.0)				(100.0)			
7.0～	67					67	2	26	95
(%)	(100.0)					(100.0)			
8.0～	126					126	1	33	160
(%)	(100.0)					(100.0)			
合計	137,894	1,487	73	3		139,457	502	33,595	173,554
(%)	(98.9)	(1.1)	(0.1)	(0.0)		(100.0)			
記載なし	2,615	18	2			2,635	17	2,003	4,655
(%)	(99.2)	(0.7)	(0.1)			(100.0)			
総計	140,509	1,505	75	3		142,092	519	35,598	178,209
(%)	(98.9)	(1.1)	(0.1)	(0.0)		(100.0)			
平均	2.48	2.59	2.45	2.10		2.48	2.31	2.50	2.49
標準偏差	1.05	0.95	0.89	0.56		1.05	1.53	1.06	1.05

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査