

表 1816 腎移植の回数 体重減少率(%)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

体重減少率	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
1.0未満	2,948	14	1			2,963	20	713	3,696
(%)	(99.5)	(0.5)	(0.0)			(100.0)			
1.0～	5,287	42	3			5,332	22	1,255	6,609
(%)	(99.2)	(0.8)	(0.1)			(100.0)			
2.0～	13,467	120	2			13,589	55	3,277	16,921
(%)	(99.1)	(0.9)	(0.0)			(100.0)			
3.0～	27,566	253	12			27,831	121	6,632	34,584
(%)	(99.0)	(0.9)	(0.0)			(100.0)			
4.0～	36,884	381	19	2		37,286	125	9,004	46,415
(%)	(98.9)	(1.0)	(0.1)	(0.0)		(100.0)			
5.0～	28,854	325	23			29,202	92	6,995	36,289
(%)	(98.8)	(1.1)	(0.1)			(100.0)			
6.0～	14,545	219	11	1		14,776	51	3,600	18,427
(%)	(98.4)	(1.5)	(0.1)	(0.0)		(100.0)			
7.0～	5,609	90	1			5,700	7	1,401	7,108
(%)	(98.4)	(1.6)	(0.0)			(100.0)			
8.0～	1,727	33	1			1,761	6	428	2,195
(%)	(98.1)	(1.9)	(0.1)			(100.0)			
9.0～	928	10				938	1	266	1,205
(%)	(98.9)	(1.1)				(100.0)			
合計	137,815	1,487	73	3		139,378	500	33,571	173,449
(%)	(98.9)	(1.1)	(0.1)	(0.0)		(100.0)			
記載なし	2,694	18	2			2,714	19	2,027	4,760
(%)	(99.3)	(0.7)	(0.1)			(100.0)			
総計	140,509	1,505	75	3		142,092	519	35,598	178,209
(%)	(98.9)	(1.1)	(0.1)	(0.0)		(100.0)			
平均	4.53	4.86	4.74	5.12		4.53	4.22	4.55	4.53
標準偏差	1.65	1.60	1.46	1.01		1.65	1.61	1.66	1.65

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査