

表 1817 腎移植の回数 Body Mass Index別（施設血液透析患者 透析回数週3回）

Body Mass Index	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
14未満	1,072	9				1,081	4	226	1,311
(%)	(99.2)	(0.8)				(100.0)			
14～	6,231	66	7	2		6,306	21	1,279	7,606
(%)	(98.8)	(1.0)	(0.1)	(0.0)		(100.0)			
16～	19,907	297	15	2		20,221	71	4,188	24,480
(%)	(98.4)	(1.5)	(0.1)	(0.0)		(100.0)			
18～	34,980	458	23			35,461	103	7,592	43,156
(%)	(98.6)	(1.3)	(0.1)			(100.0)			
20～	38,705	386	21	1		39,113	118	8,449	47,680
(%)	(99.0)	(1.0)	(0.1)	(0.0)		(100.0)			
22～	29,853	221	15			30,089	110	6,611	36,810
(%)	(99.2)	(0.7)	(0.0)			(100.0)			
24～	17,787	109	1			17,897	62	3,954	21,913
(%)	(99.4)	(0.6)	(0.0)			(100.0)			
26～	9,436	56				9,492	21	2,175	11,688
(%)	(99.4)	(0.6)				(100.0)			
28～	4,861	24	1			4,886	12	1,093	5,991
(%)	(99.5)	(0.5)	(0.0)			(100.0)			
30～	4,203	22	1			4,226	14	924	5,164
(%)	(99.5)	(0.5)	(0.0)			(100.0)			
合計	167,035	1,648	84	5		168,772	536	36,491	205,799
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
記載なし	13,736	116	4			13,856	151	9,165	23,172
(%)	(99.1)	(0.8)	(0.0)			(100.0)			
総計	180,771	1,764	88	5		182,628	687	45,656	228,971
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	21.46	20.45	19.80	17.29		21.45	21.28	21.53	21.46
標準偏差	3.66	3.25	3.12	2.52		3.66	3.53	3.66	3.66

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査