

表 1818 腎移植の回数 透析前BUN(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前BUN	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
20未満	389	2				391	1	87	479
(%)	(99.5)	(0.5)				(100.0)			
20～	1,974	9	1			1,984	16	480	2,480
(%)	(99.5)	(0.5)	(0.1)			(100.0)			
30～	8,188	40	1			8,229	40	2,089	10,358
(%)	(99.5)	(0.5)	(0.0)			(100.0)			
40～	21,364	131	10	1		21,506	90	5,215	26,811
(%)	(99.3)	(0.6)	(0.0)	(0.0)		(100.0)			
50～	33,989	294	15	1		34,299	113	8,247	42,659
(%)	(99.1)	(0.9)	(0.0)	(0.0)		(100.0)			
60～	33,972	399	18			34,389	104	8,150	42,643
(%)	(98.8)	(1.2)	(0.1)			(100.0)			
70～	22,532	334	18	1		22,885	69	5,368	28,322
(%)	(98.5)	(1.5)	(0.1)	(0.0)		(100.0)			
80～	10,420	155	7			10,582	53	2,577	13,212
(%)	(98.5)	(1.5)	(0.1)			(100.0)			
90～	3,798	82	3			3,883	10	910	4,803
(%)	(97.8)	(2.1)	(0.1)			(100.0)			
100～	1,073	30	2			1,105	6	237	1,348
(%)	(97.1)	(2.7)	(0.2)			(100.0)			
110～	323	6				329		65	394
(%)	(98.2)	(1.8)				(100.0)			
120～	163	3				166	1	37	204
(%)	(98.2)	(1.8)				(100.0)			
合計	138,185	1,485	75	3		139,748	503	33,462	173,713
(%)	(98.9)	(1.1)	(0.1)	(0.0)		(100.0)			
記載なし	2,324	20				2,344	16	2,136	4,496
(%)	(99.1)	(0.9)				(100.0)			
総計	140,509	1,505	75	3		142,092	519	35,598	178,209
(%)	(98.9)	(1.1)	(0.1)	(0.0)		(100.0)			
平均	60.93	66.72	65.61	59.67		60.99	59.54	60.76	60.94
標準偏差	15.64	15.56	15.10	15.57		15.65	16.93	15.56	15.64

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査