

表 1826 腎移植の回数 透析前補正カルシウム濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

透析前補正 カルシウム濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
7.0未満	693	20	1			714	1	159	874
(%)	(97.1)	(2.8)	(0.1)			(100.0)			
7.0～	1,017	21	1			1,039		227	1,266
(%)	(97.9)	(2.0)	(0.1)			(100.0)			
7.5～	3,869	57	3			3,929	10	1,032	4,971
(%)	(98.5)	(1.5)	(0.1)			(100.0)			
8.0～	12,850	120	8			12,978	32	3,280	16,290
(%)	(99.0)	(0.9)	(0.1)			(100.0)			
8.4～	3,762	27	1	1		3,791	7	878	4,676
(%)	(99.2)	(0.7)	(0.0)	(0.0)		(100.0)			
8.5～	42,814	382	15	1		43,212	158	10,532	53,902
(%)	(99.1)	(0.9)	(0.0)	(0.0)		(100.0)			
9.0～	52,280	489	28	2		52,799	218	12,219	65,236
(%)	(99.0)	(0.9)	(0.1)	(0.0)		(100.0)			
9.5～	34,019	358	17			34,394	101	7,909	42,404
(%)	(98.9)	(1.0)	(0.0)			(100.0)			
10.0～	4,234	41	3			4,278	9	940	5,227
(%)	(99.0)	(1.0)	(0.1)			(100.0)			
10.1～	10,684	112	5	1		10,802	37	2,438	13,277
(%)	(98.9)	(1.0)	(0.0)	(0.0)		(100.0)			
10.5～	5,125	68	3			5,196	18	1,239	6,453
(%)	(98.6)	(1.3)	(0.1)			(100.0)			
11.0～	1,749	14				1,763	8	394	2,165
(%)	(99.2)	(0.8)				(100.0)			
11.5～	625	5	2			632	7	130	769
(%)	(98.9)	(0.8)	(0.3)			(100.0)			
12.0～	444	1				445	4	90	539
(%)	(99.8)	(0.2)				(100.0)			
合計	174,165	1,715	87	5		175,972	610	41,467	218,049
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
記載なし	6,606	49	1			6,656	77	4,189	10,922
(%)	(99.2)	(0.7)	(0.0)			(100.0)			
総計	180,771	1,764	88	5		182,628	687	45,656	228,971
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	9.19	9.17	9.19	9.22		9.19	9.26	9.17	9.18
標準偏差	0.74	0.80	0.84	0.63		0.74	0.73	0.74	0.74

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査