

表 1827 腎移植の回数 透析前リン濃度(mg/dL)別 (施設血液透析患者 透析回数週3回)

透析前リン濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
3未満	7,206	48	4			7,258	48	1,655	8,961
(%)	(99.3)	(0.7)	(0.1)			(100.0)			
3.0～	8,777	57	4	1		8,839	36	2,105	10,980
(%)	(99.3)	(0.6)	(0.0)	(0.0)		(100.0)			
3.5～	15,277	99	4			15,380	68	3,570	19,018
(%)	(99.3)	(0.6)	(0.0)			(100.0)			
4.0～	22,688	182	8			22,878	86	5,228	28,192
(%)	(99.2)	(0.8)	(0.0)			(100.0)			
4.5～	26,859	237	16	2		27,114	99	6,255	33,468
(%)	(99.1)	(0.9)	(0.1)	(0.0)		(100.0)			
5.0～	26,813	241	19			27,073	105	6,631	33,809
(%)	(99.0)	(0.9)	(0.1)			(100.0)			
5.5～	23,196	258	7	1		23,462	78	5,472	29,012
(%)	(98.9)	(1.1)	(0.0)	(0.0)		(100.0)			
6.0～	3,881	44				3,925	9	994	4,928
(%)	(98.9)	(1.1)				(100.0)			
6.1～	13,127	170	3			13,300	44	3,161	16,505
(%)	(98.7)	(1.3)	(0.0)			(100.0)			
6.5～	11,384	137	11	1		11,533	34	2,810	14,377
(%)	(98.7)	(1.2)	(0.1)	(0.0)		(100.0)			
7.0～	6,938	96	4			7,038	21	1,801	8,860
(%)	(98.6)	(1.4)	(0.1)			(100.0)			
7.5～	4,199	59	4			4,262	18	1,079	5,359
(%)	(98.5)	(1.4)	(0.1)			(100.0)			
8.0～	2,535	37	3			2,575	8	609	3,192
(%)	(98.4)	(1.4)	(0.1)			(100.0)			
8.5～	1,578	25	1			1,604	2	417	2,023
(%)	(98.4)	(1.6)	(0.1)			(100.0)			
9.0～	2,479	46				2,525	4	627	3,156
(%)	(98.2)	(1.8)				(100.0)			
合計	176,937	1,736	88	5		178,766	660	42,414	221,840
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
記載なし	3,834	28				3,862	27	3,242	7,131
(%)	(99.3)	(0.7)				(100.0)			
総計	180,771	1,764	88	5		182,628	687	45,656	228,971
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	5.19	5.57	5.32	5.00		5.19	4.94	5.23	5.20
標準偏差	1.42	1.52	1.47	1.33		1.42	1.41	1.44	1.43

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査