

表 1830 腎移植の回数 透析前ヘモグロビン濃度(g/dL)別 (施設血液透析患者 透析回数週3回)

透析前ヘモグロビン 濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
5未満	233	4				237		24	261
(%)	(98.3)	(1.7)				(100.0)			
5～	313	2				315	1	46	362
(%)	(99.4)	(0.6)				(100.0)			
6～	767	4				771	6	165	942
(%)	(99.5)	(0.5)				(100.0)			
7～	2,901	29	2	1		2,933	12	633	3,578
(%)	(98.9)	(1.0)	(0.1)	(0.0)		(100.0)			
8～	9,558	78	3			9,639	45	2,042	11,726
(%)	(99.2)	(0.8)	(0.0)			(100.0)			
9～	29,745	269	14			30,028	100	6,910	37,038
(%)	(99.1)	(0.9)	(0.0)			(100.0)			
10～	60,165	536	29	1		60,731	214	14,272	75,217
(%)	(99.1)	(0.9)	(0.0)	(0.0)		(100.0)			
11～	48,960	501	32	2		49,495	182	11,802	61,479
(%)	(98.9)	(1.0)	(0.1)	(0.0)		(100.0)			
12～	17,722	232	6	1		17,961	81	4,328	22,370
(%)	(98.7)	(1.3)	(0.0)	(0.0)		(100.0)			
13～	4,418	62	1			4,481	20	1,036	5,537
(%)	(98.6)	(1.4)	(0.0)			(100.0)			
14～	1,569	15	1			1,585	6	342	1,933
(%)	(99.0)	(0.9)	(0.1)			(100.0)			
合計	176,351	1,732	88	5		178,176	667	41,600	220,443
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
記載なし	4,420	32				4,452	20	4,056	8,528
(%)	(99.3)	(0.7)				(100.0)			
総計	180,771	1,764	88	5		182,628	687	45,656	228,971
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	10.68	10.82	10.74	10.82		10.68	10.71	10.71	10.69
標準偏差	1.28	1.31	1.10	2.00		1.28	1.34	1.24	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査