

表 1840 腎移植の回数 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
慢性糸球体腎炎	11,634	291	15	1		11,941	52	3,083	15,076
(%)	(97.4)	(2.4)	(0.1)	(0.0)		(100.0)			
慢性腎盂腎炎	340	4	1			345		77	422
(%)	(98.6)	(1.2)	(0.3)			(100.0)			
急速進行性糸球体腎炎	221	1				222		62	284
(%)	(99.5)	(0.5)				(100.0)			
妊娠腎/妊娠中毒症	264	5				269	2	88	359
(%)	(98.1)	(1.9)				(100.0)			
その他の分類不能の腎炎	150	5	1			156		39	195
(%)	(96.2)	(3.2)	(0.6)			(100.0)			
多発性嚢胞腎	1,153	6	1			1,160	6	298	1,464
(%)	(99.4)	(0.5)	(0.1)			(100.0)			
腎硬化症	2,287	9	1			2,297	16	573	2,886
(%)	(99.6)	(0.4)	(0.0)			(100.0)			
悪性高血圧	244	1				245	2	65	312
(%)	(99.6)	(0.4)				(100.0)			
糖尿病性腎症	10,575	18	2			10,595	51	2,527	13,173
(%)	(99.8)	(0.2)	(0.0)			(100.0)			
SLE腎炎	301	1				302	1	60	363
(%)	(99.7)	(0.3)				(100.0)			
アミロイド腎	48					48		11	59
(%)	(100.0)					(100.0)			
痛風腎	117	1	1			119	2	33	154
(%)	(98.3)	(0.8)	(0.8)			(100.0)			
先天性代謝異常による腎不全	29	4				33		11	44
(%)	(87.9)	(12.1)				(100.0)			
腎・尿路結核	19					19		3	22
(%)	(100.0)					(100.0)			
腎・尿路結石	52					52	1	16	69
(%)	(100.0)					(100.0)			
腎・尿路腫瘍	72					72		13	85
(%)	(100.0)					(100.0)			
閉塞性尿路障害	57	1				58		16	74
(%)	(98.3)	(1.7)				(100.0)			
骨髄腫	16					16		3	19
(%)	(100.0)					(100.0)			
腎形成不全	69	7				76	1	24	101
(%)	(90.8)	(9.2)				(100.0)			
不明	2,680	42	2			2,724	9	747	3,480
(%)	(98.4)	(1.5)	(0.1)			(100.0)			
再導入	104	240	11	1		356		60	416
(%)	(29.2)	(67.4)	(3.1)	(0.3)		(100.0)			
その他	647	20	1			668	3	218	889
(%)	(96.9)	(3.0)	(0.1)			(100.0)			
合計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
記載なし									
(%)									
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査