

表 1844 腎移植の回数 身長(cm)別（血液透析濾過患者 透析回数週3回）

身長	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
130未満	32					32	1	5	38
(%)	(100.0)					(100.0)			
130～	48	1	1			50	1	19	70
(%)	(96.0)	(2.0)	(2.0)			(100.0)			
135～	245	2	1			248	1	40	289
(%)	(98.8)	(0.8)	(0.4)			(100.0)			
140～	936	9		1		946	9	208	1,163
(%)	(98.9)	(1.0)		(0.1)		(100.0)			
145～	2,372	54	1			2,427	10	558	2,995
(%)	(97.7)	(2.2)	(0.0)			(100.0)			
150～	4,231	69	3			4,303	25	884	5,212
(%)	(98.3)	(1.6)	(0.1)			(100.0)			
155～	4,882	88	8			4,978	27	1,074	6,079
(%)	(98.1)	(1.8)	(0.2)			(100.0)			
160～	5,768	137	7	1		5,913	25	1,239	7,177
(%)	(97.5)	(2.3)	(0.1)	(0.0)		(100.0)			
165～	5,426	137	6			5,569	14	1,139	6,722
(%)	(97.4)	(2.5)	(0.1)			(100.0)			
170～	3,853	87	3			3,943	20	822	4,785
(%)	(97.7)	(2.2)	(0.1)			(100.0)			
175～	1,434	35	4			1,473	6	307	1,786
(%)	(97.4)	(2.4)	(0.3)			(100.0)			
180～	405	13				418	1	74	493
(%)	(96.9)	(3.1)				(100.0)			
185～	57	3	1			61		16	77
(%)	(93.4)	(4.9)	(1.6)			(100.0)			
190～	8					8		1	9
(%)	(100.0)					(100.0)			
合計	29,697	635	35	2		30,369	140	6,386	36,895
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
記載なし	1,382	21	1			1,404	6	1,641	3,051
(%)	(98.4)	(1.5)	(0.1)			(100.0)			
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
平均	160.75	162.27	162.40	152.00		160.78	158.85	160.58	160.74
標準偏差	9.55	9.22	11.25	11.31		9.54	10.31	9.48	9.54

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査