

表 1845 腎移植の回数 透析後体重(kg)別（血液透析濾過患者 透析回数週3回）

透析後体重	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
20未満	8					8	1	3	12
(%)	(100.0)					(100.0)			
20～	58					58		18	76
(%)	(100.0)					(100.0)			
30～	2,078	48	2			2,128	14	540	2,682
(%)	(97.7)	(2.3)	(0.1)			(100.0)			
40～	7,875	189	11	1		8,076	32	1,897	10,005
(%)	(97.5)	(2.3)	(0.1)	(0.0)		(100.0)			
50～	9,790	230	13	1		10,034	47	2,409	12,490
(%)	(97.6)	(2.3)	(0.1)	(0.0)		(100.0)			
60～	6,628	122	5			6,755	22	1,588	8,365
(%)	(98.1)	(1.8)	(0.1)			(100.0)			
70～	2,723	47	2			2,772	10	641	3,423
(%)	(98.2)	(1.7)	(0.1)			(100.0)			
80～	1,029	10	1			1,040	6	227	1,273
(%)	(98.9)	(1.0)	(0.1)			(100.0)			
90～	359	3				362	1	92	455
(%)	(99.2)	(0.8)				(100.0)			
100～	108	1				109	1	32	142
(%)	(99.1)	(0.9)				(100.0)			
110～	49					49		12	61
(%)	(100.0)					(100.0)			
120～	29					29		8	37
(%)	(100.0)					(100.0)			
合計	30,734	650	34	2		31,420	134	7,467	39,021
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
記載なし	345	6	2			353	12	560	925
(%)	(97.7)	(1.7)	(0.6)			(100.0)			
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
平均	56.72	54.46	54.03	48.65		56.67	55.46	56.48	56.63
標準偏差	13.03	11.07	10.84	10.68		13.00	14.21	13.15	13.03

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査