

表 1846 腎移植の回数 体重減少量(kg)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

体重減少量	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
0.0未満	79	3				82		15	97
(%)	(96.3)	(3.7)				(100.0)			
0.0～	1,054	20		1		1,075	4	241	1,320
(%)	(98.0)	(1.9)		(0.1)		(100.0)			
1.0～	5,801	123	5	1		5,930	18	1,452	7,400
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
2.0～	11,483	287	19			11,789	60	2,805	14,654
(%)	(97.4)	(2.4)	(0.2)			(100.0)			
3.0～	6,286	139	5			6,430	21	1,563	8,014
(%)	(97.8)	(2.2)	(0.1)			(100.0)			
4.0～	1,716	33	2			1,751	10	430	2,191
(%)	(98.0)	(1.9)	(0.1)			(100.0)			
5.0～	301	7	1			309		91	400
(%)	(97.4)	(2.3)	(0.3)			(100.0)			
6.0～	49					49		10	59
(%)	(100.0)					(100.0)			
7.0～	11	1				12		4	16
(%)	(91.7)	(8.3)				(100.0)			
8.0～	23					23		7	30
(%)	(100.0)					(100.0)			
合計	26,803	613	32	2		27,450	113	6,618	34,181
(%)	(97.6)	(2.2)	(0.1)	(0.0)		(100.0)			
記載なし	300	4	2			306	12	472	790
(%)	(98.0)	(1.3)	(0.7)			(100.0)			
総計	27,103	617	34	2		27,756	125	7,090	34,971
(%)	(97.6)	(2.2)	(0.1)	(0.0)		(100.0)			
平均	2.56	2.58	2.68	1.05		2.56	2.64	2.58	2.57
標準偏差	1.01	0.94	0.85	0.21		1.01	0.84	1.10	1.03

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査