

表 1847 腎移植の回数 体重減少率(%)別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

体重減少率	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
1.0未満	482	10				492	1	94	587
(%)	(98.0)	(2.0)				(100.0)			
1.0～	851	16		1		868	3	220	1,091
(%)	(98.0)	(1.8)		(0.1)		(100.0)			
2.0～	2,399	44	1	1		2,445	3	582	3,030
(%)	(98.1)	(1.8)	(0.0)	(0.0)		(100.0)			
3.0～	5,452	114	6			5,572	16	1,305	6,893
(%)	(97.8)	(2.0)	(0.1)			(100.0)			
4.0～	7,360	161	10			7,531	37	1,809	9,377
(%)	(97.7)	(2.1)	(0.1)			(100.0)			
5.0～	5,780	141	8			5,929	33	1,462	7,424
(%)	(97.5)	(2.4)	(0.1)			(100.0)			
6.0～	2,879	72	3			2,954	16	755	3,725
(%)	(97.5)	(2.4)	(0.1)			(100.0)			
7.0～	1,076	43	3			1,122	4	282	1,408
(%)	(95.9)	(3.8)	(0.3)			(100.0)			
8.0～	356	7	1			364		72	436
(%)	(97.8)	(1.9)	(0.3)			(100.0)			
9.0～	154	5				159		31	190
(%)	(96.9)	(3.1)				(100.0)			
合計	26,789	613	32	2		27,436	113	6,612	34,161
(%)	(97.6)	(2.2)	(0.1)	(0.0)		(100.0)			
記載なし	314	4	2			320	12	478	810
(%)	(98.1)	(1.3)	(0.6)			(100.0)			
総計	27,103	617	34	2		27,756	125	7,090	34,971
(%)	(97.6)	(2.2)	(0.1)	(0.0)		(100.0)			
平均	4.58	4.79	5.09	2.26		4.59	4.88	4.61	4.59
標準偏差	1.59	1.63	1.39	0.93		1.59	1.23	1.55	1.59

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査