

表 1848 腎移植の回数 Body Mass Index別（血液透析濾過患者 透析回数週3回）

Body Mass Index	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
14未満	118	4				122		27	149
(%)	(96.7)	(3.3)				(100.0)			
14～	812	25	1			838	4	204	1,046
(%)	(96.9)	(3.0)	(0.1)			(100.0)			
16～	3,173	94	8			3,275	12	711	3,998
(%)	(96.9)	(2.9)	(0.2)			(100.0)			
18～	6,139	175	4	1		6,319	29	1,266	7,614
(%)	(97.2)	(2.8)	(0.1)	(0.0)		(100.0)			
20～	6,793	152	12	1		6,958	23	1,481	8,462
(%)	(97.6)	(2.2)	(0.2)	(0.0)		(100.0)			
22～	5,311	101	6			5,418	26	1,127	6,571
(%)	(98.0)	(1.9)	(0.1)			(100.0)			
24～	3,359	40	2			3,401	11	676	4,088
(%)	(98.8)	(1.2)	(0.1)			(100.0)			
26～	1,779	20				1,799	12	386	2,197
(%)	(98.9)	(1.1)				(100.0)			
28～	970	13				983	5	189	1,177
(%)	(98.7)	(1.3)				(100.0)			
30～	851	6				857	4	156	1,017
(%)	(99.3)	(0.7)				(100.0)			
合計	29,305	630	33	2		29,970	126	6,223	36,319
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
記載なし	1,774	26	3			1,803	20	1,804	3,627
(%)	(98.4)	(1.4)	(0.2)			(100.0)			
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
平均	21.71	20.60	20.12	20.89		21.69	22.02	21.58	21.67
標準偏差	3.66	3.20	2.71	1.51		3.65	3.97	3.62	3.65

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査