

表 1849 腎移植の回数 透析前BUN(mg/dL)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

透析前BUN	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
20未満	29	1				30		9	39
(%)	(96.7)	(3.3)				(100.0)			
20～	256	3				259	4	46	309
(%)	(98.8)	(1.2)				(100.0)			
30～	1,318	17				1,335	8	310	1,653
(%)	(98.7)	(1.3)				(100.0)			
40～	3,815	52	3			3,870	17	898	4,785
(%)	(98.6)	(1.3)	(0.1)			(100.0)			
50～	6,572	128	7	1		6,708	31	1,579	8,318
(%)	(98.0)	(1.9)	(0.1)	(0.0)		(100.0)			
60～	6,905	182	6			7,093	26	1,685	8,804
(%)	(97.3)	(2.6)	(0.1)			(100.0)			
70～	4,697	133	8			4,838	22	1,226	6,086
(%)	(97.1)	(2.7)	(0.2)			(100.0)			
80～	2,227	66	4			2,297	5	529	2,831
(%)	(97.0)	(2.9)	(0.2)			(100.0)			
90～	807	25	2	1		835	2	214	1,051
(%)	(96.6)	(3.0)	(0.2)	(0.1)		(100.0)			
100～	210	3	1			214		65	279
(%)	(98.1)	(1.4)	(0.5)			(100.0)			
110～	58	2	1			61		13	74
(%)	(95.1)	(3.3)	(1.6)			(100.0)			
120～	16	2				18		2	20
(%)	(88.9)	(11.1)				(100.0)			
合計	26,910	614	32	2		27,558	115	6,576	34,249
(%)	(97.6)	(2.2)	(0.1)	(0.0)		(100.0)			
記載なし	193	3	2			198	10	514	722
(%)	(97.5)	(1.5)	(1.0)			(100.0)			
総計	27,103	617	34	2		27,756	125	7,090	34,971
(%)	(97.6)	(2.2)	(0.1)	(0.0)		(100.0)			
平均	62.08	65.87	69.63	75.50		62.18	58.83	62.52	62.23
標準偏差	15.09	15.10	17.70	23.33		15.10	14.64	15.00	15.08

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査