

表 1856 腎移植の回数 透析前CRP濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

透析前CRP濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
0.2未満	16,480	364	18	1		16,863	56	3,785	20,704
(%)	(97.7)	(2.2)	(0.1)	(0.0)		(100.0)			
0.2～	5,172	106	7			5,285	24	1,168	6,477
(%)	(97.9)	(2.0)	(0.1)			(100.0)			
0.5～	2,504	42	3			2,549	10	614	3,173
(%)	(98.2)	(1.6)	(0.1)			(100.0)			
1.0～	1,588	31	1			1,620	4	379	2,003
(%)	(98.0)	(1.9)	(0.1)			(100.0)			
2.0～	918	14	3			935	1	198	1,134
(%)	(98.2)	(1.5)	(0.3)			(100.0)			
4.0～	317	9				326		81	407
(%)	(97.2)	(2.8)				(100.0)			
6.0～	138	1				139	1	31	171
(%)	(99.3)	(0.7)				(100.0)			
8.0～	64			1		65		29	94
(%)	(98.5)			(1.5)		(100.0)			
10.0～	83	4				87		21	108
(%)	(95.4)	(4.6)				(100.0)			
15.0～	63	1				64		20	84
(%)	(98.4)	(1.6)				(100.0)			
合計	27,327	572	32	2		27,933	96	6,326	34,355
(%)	(97.8)	(2.0)	(0.1)	(0.0)		(100.0)			
記載なし	3,752	84	4			3,840	50	1,701	5,591
(%)	(97.7)	(2.2)	(0.1)			(100.0)			
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
平均	0.54	0.49	0.48	4.17		0.54	0.33	0.60	0.55
標準偏差	1.69	1.37	0.79	5.76		1.68	0.73	2.09	1.76

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査