

表 1857 腎移植の回数 透析前補正カルシウム濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

透析前補正 カルシウム濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
7.0未満	138	3				141		19	160
(%)	(97.9)	(2.1)				(100.0)			
7.0～	215	10				225		34	259
(%)	(95.6)	(4.4)				(100.0)			
7.5～	749	17	2			768	1	182	951
(%)	(97.5)	(2.2)	(0.3)			(100.0)			
8.0～	2,271	54				2,325	4	539	2,868
(%)	(97.7)	(2.3)				(100.0)			
8.4～	660	12	1			673	2	143	818
(%)	(98.1)	(1.8)	(0.1)			(100.0)			
8.5～	7,468	114	12	1		7,595	33	1,673	9,301
(%)	(98.3)	(1.5)	(0.2)	(0.0)		(100.0)			
9.0～	9,181	190	7			9,378	28	2,109	11,515
(%)	(97.9)	(2.0)	(0.1)			(100.0)			
9.5～	5,926	140	7	1		6,074	20	1,518	7,612
(%)	(97.6)	(2.3)	(0.1)	(0.0)		(100.0)			
10.0～	700	16				716	5	182	903
(%)	(97.8)	(2.2)				(100.0)			
10.1～	1,913	48	5			1,966	6	480	2,452
(%)	(97.3)	(2.4)	(0.3)			(100.0)			
10.5～	923	24				947	3	220	1,170
(%)	(97.5)	(2.5)				(100.0)			
11.0～	296	10				306		77	383
(%)	(96.7)	(3.3)				(100.0)			
11.5～	104	4				108		24	132
(%)	(96.3)	(3.7)				(100.0)			
12.0～	58	3				61		10	71
(%)	(95.1)	(4.9)				(100.0)			
合計	30,602	645	34	2		31,283	102	7,210	38,595
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
記載なし	477	11	2			490	44	817	1,351
(%)	(97.3)	(2.2)	(0.4)			(100.0)			
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
平均	9.18	9.24	9.17	9.30		9.18	9.21	9.21	9.19
標準偏差	0.75	0.83	0.66	0.71		0.75	0.61	0.73	0.75

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査