

表 1858 腎移植の回数 透析前リン濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

透析前リン濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
3未満	914	8	1			923	4	206	1,133
(%)	(99.0)	(0.9)	(0.1)			(100.0)			
3.0～	1,263	16				1,279	9	286	1,574
(%)	(98.7)	(1.3)				(100.0)			
3.5～	2,278	35	2			2,315	11	523	2,849
(%)	(98.4)	(1.5)	(0.1)			(100.0)			
4.0～	3,608	62	3	1		3,674	14	877	4,565
(%)	(98.2)	(1.7)	(0.1)	(0.0)		(100.0)			
4.5～	4,525	95	6			4,626	22	1,064	5,712
(%)	(97.8)	(2.1)	(0.1)			(100.0)			
5.0～	4,860	114	5			4,979	18	1,137	6,134
(%)	(97.6)	(2.3)	(0.1)			(100.0)			
5.5～	4,180	105	6			4,291	18	990	5,299
(%)	(97.4)	(2.4)	(0.1)			(100.0)			
6.0～	739	14				753	2	195	950
(%)	(98.1)	(1.9)				(100.0)			
6.1～	2,559	47	2			2,608	12	624	3,244
(%)	(98.1)	(1.8)	(0.1)			(100.0)			
6.5～	2,217	67	2			2,286	10	510	2,806
(%)	(97.0)	(2.9)	(0.1)			(100.0)			
7.0～	1,397	30	3			1,430	7	338	1,775
(%)	(97.7)	(2.1)	(0.2)			(100.0)			
7.5～	881	16	2	1		900	2	202	1,104
(%)	(97.9)	(1.8)	(0.2)	(0.1)		(100.0)			
8.0～	509	17				526	3	122	651
(%)	(96.8)	(3.2)				(100.0)			
8.5～	344	10				354		69	423
(%)	(97.2)	(2.8)				(100.0)			
9.0～	522	16	2			540	2	140	682
(%)	(96.7)	(3.0)	(0.4)			(100.0)			
合計	30,796	652	34	2		31,484	134	7,283	38,901
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
記載なし	283	4	2			289	12	744	1,045
(%)	(97.9)	(1.4)	(0.7)			(100.0)			
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
平均	5.35	5.59	5.68	5.80		5.35	5.24	5.37	5.36
標準偏差	1.42	1.38	1.53	2.40		1.42	1.40	1.43	1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査