

表 1861 腎移植の回数 透析前ヘモグロビン濃度(g/dL)別 (血液透析濾過患者 透析回数週3回)

透析前ヘモグロビン 濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
5未満	22		1			23		3	26
(%)	(95.7)		(4.3)			(100.0)			
5～	30					30		1	31
(%)	(100.0)					(100.0)			
6～	93	3				96	1	17	114
(%)	(96.9)	(3.1)				(100.0)			
7～	373	8				381	2	62	445
(%)	(97.9)	(2.1)				(100.0)			
8～	1,328	32	1			1,361	13	263	1,637
(%)	(97.6)	(2.4)	(0.1)			(100.0)			
9～	4,647	79	4			4,730	32	1,088	5,850
(%)	(98.2)	(1.7)	(0.1)			(100.0)			
10～	10,171	208	12	1		10,392	50	2,505	12,947
(%)	(97.9)	(2.0)	(0.1)	(0.0)		(100.0)			
11～	9,111	199	9	1		9,320	28	2,229	11,577
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
12～	3,472	84	6			3,562	6	814	4,382
(%)	(97.5)	(2.4)	(0.2)			(100.0)			
13～	932	27	1			960	2	223	1,185
(%)	(97.1)	(2.8)	(0.1)			(100.0)			
14～	420	9				429	1	93	523
(%)	(97.9)	(2.1)				(100.0)			
合計	30,599	649	34	2		31,284	135	7,298	38,717
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
記載なし	480	7	2			489	11	729	1,229
(%)	(98.2)	(1.4)	(0.4)			(100.0)			
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
平均	10.83	10.92	10.80	10.85		10.83	10.33	10.87	10.84
標準偏差	1.28	1.31	1.65	0.64		1.28	1.20	1.20	1.26

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査