

表 1863 腎移植の回数 HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

HDL-C濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
20未満	64	1				65		16	81
(%)	(98.5)	(1.5)				(100.0)			
20～	1,475	20	2			1,497	7	370	1,874
(%)	(98.5)	(1.3)	(0.1)			(100.0)			
30～	4,928	87	3			5,018	20	1,093	6,131
(%)	(98.2)	(1.7)	(0.1)			(100.0)			
40～	5,807	121	7			5,935	29	1,396	7,360
(%)	(97.8)	(2.0)	(0.1)			(100.0)			
50～	4,694	111	6	2		4,813	11	1,147	5,971
(%)	(97.5)	(2.3)	(0.1)	(0.0)		(100.0)			
60～	3,082	63	2			3,147	8	702	3,857
(%)	(97.9)	(2.0)	(0.1)			(100.0)			
70～	1,567	42	1			1,610	6	407	2,023
(%)	(97.3)	(2.6)	(0.1)			(100.0)			
80～	782	26	2			810	3	200	1,013
(%)	(96.5)	(3.2)	(0.2)			(100.0)			
90～	322	8				330	1	100	431
(%)	(97.6)	(2.4)				(100.0)			
100～	277	13				290	1	92	383
(%)	(95.5)	(4.5)				(100.0)			
合計	22,998	492	23	2		23,515	86	5,523	29,124
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
記載なし	8,081	164	13			8,258	60	2,504	10,822
(%)	(97.9)	(2.0)	(0.2)			(100.0)			
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
平均	50.66	54.22	50.96	52.00		50.73	48.59	51.41	50.85
標準偏差	17.02	18.62	15.73	2.83		17.06	16.86	17.71	17.18

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査