

表 1864 腎移植の回数 Non-HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

Non-HDL-C濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
50未満	418	12				430	1	98	529
(%)	(97.2)	(2.8)				(100.0)			
50～	1,835	40	3			1,878	1	524	2,403
(%)	(97.7)	(2.1)	(0.2)			(100.0)			
70～	4,060	78	7	1		4,146	2	990	5,138
(%)	(97.9)	(1.9)	(0.2)	(0.0)		(100.0)			
90～	4,947	113	4			5,064	2	1,199	6,265
(%)	(97.7)	(2.2)	(0.1)			(100.0)			
110～	4,048	97	4			4,149	3	1,010	5,162
(%)	(97.6)	(2.3)	(0.1)			(100.0)			
130～	2,519	47	1			2,567		582	3,149
(%)	(98.1)	(1.8)	(0.0)			(100.0)			
150～	1,264	32	2	1		1,299		298	1,597
(%)	(97.3)	(2.5)	(0.2)	(0.1)		(100.0)			
170～	550	7				557		125	682
(%)	(98.7)	(1.3)				(100.0)			
190～	200	3				203		59	262
(%)	(98.5)	(1.5)				(100.0)			
210～	102	3				105		36	141
(%)	(97.1)	(2.9)				(100.0)			
合計	19,943	432	21	2		20,398	9	4,921	25,328
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
記載なし	11,136	224	15			11,375	137	3,106	14,618
(%)	(97.9)	(2.0)	(0.1)			(100.0)			
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
平均	107.22	107.31	99.62	113.50		107.22	90.67	106.74	107.12
標準偏差	33.21	34.41	29.56	57.28		33.23	26.94	33.95	33.37

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査