

表 1867 腎移植の回数 透析前脈拍(拍/分)別 (血液透析濾過患者 透析回数週3回)

透析前脈拍	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
40未満	13					13		4	17
(%)	(100.0)					(100.0)			
40～	253	7				260	1	34	295
(%)	(97.3)	(2.7)				(100.0)			
50～	2,104	43	1			2,148	9	475	2,632
(%)	(98.0)	(2.0)	(0.0)			(100.0)			
60～	6,719	137	13			6,869	39	1,582	8,490
(%)	(97.8)	(2.0)	(0.2)			(100.0)			
70～	8,346	191	11	2		8,550	46	1,949	10,545
(%)	(97.6)	(2.2)	(0.1)	(0.0)		(100.0)			
80～	5,551	114	3			5,668	23	1,372	7,063
(%)	(97.9)	(2.0)	(0.1)			(100.0)			
90～	2,610	61	2			2,673	5	615	3,293
(%)	(97.6)	(2.3)	(0.1)			(100.0)			
100～	822	18				840	1	201	1,042
(%)	(97.9)	(2.1)				(100.0)			
110～	211	4				215		46	261
(%)	(98.1)	(1.9)				(100.0)			
120～	91					91		15	106
(%)	(100.0)					(100.0)			
合計	26,720	575	30	2		27,327	124	6,293	33,744
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
記載なし	4,359	81	6			4,446	22	1,734	6,202
(%)	(98.0)	(1.8)	(0.1)			(100.0)			
総計	31,079	656	36	2		31,773	146	8,027	39,946
(%)	(97.8)	(2.1)	(0.1)	(0.0)		(100.0)			
平均	75.39	75.52	72.33	71.50		75.39	72.35	75.63	75.43
標準偏差	13.01	12.46	10.36	0.71		12.99	10.11	12.70	12.93

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査