

表 1873 腎移植の回数 身長(cm)別（腹膜透析患者）

身長	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
130未満	56	1	1			58		4	62
(%)	(96.6)	(1.7)	(1.7)			(100.0)			
130～	10		1			11			11
(%)	(90.9)		(9.1)			(100.0)			
135～	52					52		10	62
(%)	(100.0)					(100.0)			
140～	182	1				183		44	227
(%)	(99.5)	(0.5)				(100.0)			
145～	363	2				365		105	470
(%)	(99.5)	(0.5)				(100.0)			
150～	689	5				694	1	172	867
(%)	(99.3)	(0.7)				(100.0)			
155～	844	12		1		857	3	189	1,049
(%)	(98.5)	(1.4)		(0.1)		(100.0)			
160～	1,058	7				1,065	3	262	1,330
(%)	(99.3)	(0.7)				(100.0)			
165～	966	11				977	1	234	1,212
(%)	(98.9)	(1.1)				(100.0)			
170～	724	4				728	3	173	904
(%)	(99.5)	(0.5)				(100.0)			
175～	293	4				297		69	366
(%)	(98.7)	(1.3)				(100.0)			
180～	76	1				77		10	87
(%)	(98.7)	(1.3)				(100.0)			
185～	8	1				9		4	13
(%)	(88.9)	(11.1)				(100.0)			
190～	1					1		3	4
(%)	(100.0)					(100.0)			
合計	5,322	49	2	1		5,374	11	1,279	6,664
(%)	(99.0)	(0.9)	(0.0)	(0.0)		(100.0)			
記載なし	453	7				460	12	1,805	2,277
(%)	(98.5)	(1.5)				(100.0)			
総計	5,775	56	2	1		5,834	23	3,084	8,941
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	160.49	161.45	119.00	156.00		160.48	163.36	160.83	160.56
標準偏差	11.76	13.50	15.56			11.80	7.55	10.71	11.59

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査